

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
16 June 2005 (16.06.2005)

PCT

(10) International Publication Number
WO 2005/054864 A2

(51) International Patent Classification⁷: **G01N 33/68**

(21) International Application Number:
PCT/EP2004/012785

(22) International Filing Date:
11 November 2004 (11.11.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
03027007.8 22 November 2003 (22.11.2003) EP

(71) Applicant (for all designated States except US): **BAYER HEALTHCARE AG** [DE/DE]; 51368 Leverkusen (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GOLZ, Stefan** [DE/DE]; Bückmannsmühle 46, 45326 Essen (DE). **BRÜGGEMEIER, Ulf** [DE/DE]; Leysiefen 20, 42799 Leichlingen (DE). **GEERTS, Andreas** [DE/DE]; Schuckertstr. 29, 42113 Wuppertal (DE).

(74) Common Representative: **BAYER HEALTHCARE AG**; Law and Patents, Patents and Licensing, 51368 Leverkusen (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **DIAGNOSTICS AND THERAPEUTICS FOR DISEASES ASSOCIATED WITH POTASSIUM CHANNEL, SUB-FAMILY K, MEMBER 1 (KCNK1)**

(57) Abstract: The invention provides a human KCNK1 which is associated with the cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases. The invention also provides assays for the identification of compounds useful in the treatment or prevention of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases. The invention also features compounds which bind to and/or activate or inhibit the activity of KCNK1 as well as pharmaceutical compositions comprising such compounds.



WO 2005/054864 A2

Diagnostics and Therapeutics for Diseases Associated with Potassium Channel, Subfamily K, Member 1 (KCNK1)

Technical field of the invention

The present invention is in the field of molecular biology, more particularly, the present invention relates to nucleic acid sequences and amino acid sequences of a human KCNK1 and its regulation for the treatment of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in mammals.

Background of the invention

10 Ion Channels

Ion channels are membrane-spanning proteins involved in the regulation of ion movements across biological membranes and play a crucial role in maintaining and modulating cellular homeostasis of excitable and non-excitable cells [Lesage et al. (1996b); Lesage et al. (1996a); US 6309855; WO 01/94629]. Ion channels mediate many fundamental and medically significant biological signalling processes such as secretion of hormones and neurotransmitters, generation of electric signals in the cardiovascular system and the conversion of chemical and mechanical extracellular stimuli into electrical responses in the nervous system. The broad functional diversity of ion channels is based on their specificity for ions (e.g. anions, cations, potassium, sodium, calcium, or chloride), their susceptibility to regulation and their molecular diversity. Activation and/or opening of ion channels can be voltage- or ligand-dependent or a combination of both. Examples for ligand-gated channels comprise nicotinic acetylcholine receptors, gamma-amino butyric acid (GABA) receptors, glycine receptors, and glutamate receptors. Examples for voltage-gated channels are the highly diversified families of calcium, potassium- and sodium channels. Most of the ion channel proteins are complex multisubunit macromolecules with several membrane-spanning helical regions forming the channel pore and accessory subunits with regulatory functions.

The almost ubiquitous expression of ion channels and their obvious functional importance makes them excellent targets for pharmaceutical intervention. Their accessibility to peptidic and small compound modulators is proven by many examples, such as the classical calcium antagonists verapamil, diltazem and nimodipin, the peptidic N-type calcium channel blocker family of conotoxins, locally applied anaesthetics like the sodium channel blocker lidocaine, anti-diabetic potassium channel modulators based on sulfonylurea derivatives or other pharmacologically active

compounds acting on glycine receptors, GABA receptors, acetylcholine receptors, vanilloid receptors as well as the diverse family of ionotropic glutamate receptors. In comparison to standard (non-ion channel) receptor pharmacology screening for ion channel ligands by high throughput approaches has just recently become fully available because of the specific technical challenges to measure ion channel activity on a large scale. Now that such techniques become more and more available, pharmaceutical research and development must focus on (i) expression profiling of ion channels defining their disease specific functions and (ii) the identification of selective novel drugs for therapeutic approaches to fully exploit the pharmacological potential of ion channels in order to prevent, ameliorate or correct dysfunctions or diseases with high medical need. In particular, the diseases to be addressed include, but are not limited to, infections such as bacterial, fungal, protozoan, and viral infections, particularly those caused by HIV viruses, cancers, allergies including asthma, cardiovascular diseases including acute heart failure, hypotension, hypertension, angina pectoris, myocardial infarction, hematological diseases, genito-urinary diseases including urinary incontinence and benign prostate hyperplasia, osteoporosis, and peripheral and central nervous system disorders including pain, Alzheimer's disease and Parkinson's disease (Willumsen et al., 2003 Receptors and Channels 9,3-12).

TaqMan-Technology / expression profiling

TaqMan is a recently developed technique, in which the release of a fluorescent reporter dye from a hybridisation probe in real-time during a polymerase chain reaction (PCR) is proportional to the accumulation of the PCR product. Quantification is based on the early, linear part of the reaction, and by determining the threshold cycle (CT), at which fluorescence above background is first detected.

Gene expression technologies may be useful in several areas of drug discovery and development, such as target identification, lead optimization, and identification of mechanisms of action. The TaqMan technology can be used to compare differences between expression profiles of normal tissue and diseased tissue. Expression profiling has been used in identifying genes, which are up- or downregulated in a variety of diseases. An interesting application of expression profiling is temporal monitoring of changes in gene expression during disease progression and drug treatment or in patients versus healthy individuals. The premise in this approach is that changes in pattern of gene expression in response to physiological or environmental stimuli (e.g., drugs) may serve as indirect clues about disease-causing genes or drug targets. Moreover, the effects of drugs with established efficacy on global gene expression patterns may provide a guidepost, or a genetic signature, against which a new drug candidate can be compared.

KCNK1

The nucleotide sequence of KCNK1 is accessible in the databases by the accession number NM_002245 and is given in SEQ ID NO: 1. The amino acid sequence of KCNK1 depicted in SEQ ID NO: 2.

Potassium channels are functionally important to a large number of cellular processes including maintenance of the action potential, muscle contraction, hormone secretion, osmotic regulation, and ion flow.

Lesage et al. [Lesage et al., (1996a)] cloned a member of a new class of potassium channels by computer identification of an EST with predicted sequence similarity to the P domain (a region involved in the potassium conduction pathway) of previously known channels. A cDNA, designated TWIK1 by the authors, was obtained from a human kidney library and shown to encode a predicted 336-amino acid protein. Unlike other potassium channels, TWIK1 has 4 (rather than 6) transmembrane domains and 2 (rather than 1) P domains. Two genes from *C. elegans* are related. When expressed in *Xenopus* oocytes, TWIK1 is able to direct expression of weakly inward-rectifying potassium currents. Northern blots showed that the gene is transcribed in a large number of tissues but is especially highly expressed in the brain and heart. The authors speculated that TWIK1 channels may be involved in the control of background potassium membrane conductances.

KCNK1 is published in [Lesage et al. (1996a), Lesage et al. (1996b)] and patent US 6309855 and WO0194629.

Summary of the invention

The invention relates to novel disease associations of KCNK1 polypeptides and polynucleotides. The invention also relates to novel methods of screening for therapeutic agents for the treatment of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal. The invention also relates to pharmaceutical compositions for the treatment of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising a KCNK1 polypeptide, a KCNK1 polynucleotide, or regulators of KCNK1 or modulators of KCNK1 activity. The invention further comprises methods of diagnosing cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal.

Brief Description of the Drawings

- Fig. 1 shows the nucleotide sequence of a KCNK1 polynucleotide (SEQ ID NO: 1).
Fig. 2 shows the amino acid sequence of a KCNK1 polypeptide (SEQ ID NO: 2).
Fig. 3 shows the nucleotide sequence of a primer useful for the invention (SEQ ID NO: 3).
5 Fig. 4 shows the nucleotide sequence of a primer useful for the invention (SEQ ID NO: 4).
Fig. 5 shows a nucleotide sequence useful as a probe to detect proteins of the invention (SEQ ID NO: 5).

Detailed description of the invention**Definition of terms**

- 10 An "oligonucleotide" is a stretch of nucleotide residues which has a sufficient number of bases to be used as an oligomer, amplimer or probe in a polymerase chain reaction (PCR). Oligonucleotides are prepared from genomic or cDNA sequence and are used to amplify, reveal, or confirm the presence of a similar DNA or RNA in a particular cell or tissue. Oligonucleotides or oligomers comprise portions of a DNA sequence having at least about 10 nucleotides and as many as about
15 35 nucleotides, preferably about 25 nucleotides.

- "Probes" may be derived from naturally occurring or recombinant single- or double-stranded nucleic acids or may be chemically synthesized. They are useful in detecting the presence of identical or similar sequences. Such probes may be labeled with reporter molecules using nick translation, Klenow fill-in reaction, PCR or other methods well known in the art. Nucleic acid
20 probes may be used in southern, northern or in situ hybridizations to determine whether DNA or RNA encoding a certain protein is present in a cell type, tissue, or organ.

A "fragment of a polynucleotide" is a nucleic acid that comprises all or any part of a given nucleotide molecule, the fragment having fewer nucleotides than about 6 kb, preferably fewer than about 1 kb.

- 25 "Reporter molecules" are radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents which associate with a particular nucleotide or amino acid sequence, thereby establishing the presence of a certain sequence, or allowing for the quantification of a certain sequence.

- "Chimeric" molecules may be constructed by introducing all or part of the nucleotide sequence of this invention into a vector containing additional nucleic acid sequence which might be expected to
30 change any one or several of the following KCNK1 characteristics: cellular location, distribution, ligand-binding affinities, interchain affinities, degradation/turnover rate, signaling, etc.

“Active“, with respect to a KCNK1 polypeptide, refers to those forms, fragments, or domains of a KCNK1 polypeptide which retain the biological and/or antigenic activity of a KCNK1 polypeptide.

5 “Functional activity” of ion channels refers to their ability to conduct and control ion movements across cellular membranes resulting in measurable changes in (1) the current, (2) the membrane potential, (3) the change in concentration of the transported ions and (4) any other measurable change exerted by KCNK1.

10 Depending on the type of channel, functional activity of said channel is determined by the activation state and gating properties of the channel which are influenced by parameters including but not limited to membrane potential, pH, temperature, physiological ligands, or a combination of two or more of these parameters and can be regulated by modulators such as peptides, toxins, antibodies or small compounds.

15 “Naturally occurring KCNK1 polypeptide“ refers to a polypeptide produced by cells which have not been genetically engineered and specifically contemplates various polypeptides arising from post-translational modifications of the polypeptide including but not limited to acetylation, carboxylation, glycosylation, phosphorylation, lipidation and acylation.

20 “Derivative“ refers to polypeptides which have been chemically modified by techniques such as ubiquitination, labeling (see above), pegylation (derivatization with polyethylene glycol), and chemical insertion or substitution of amino acids such as ornithine which do not normally occur in human proteins.

“Conservative amino acid substitutions“ result from replacing one amino acid with another having similar structural and/or chemical properties, such as the replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, or a threonine with a serine.

25 “Insertions“ or “deletions“ are typically in the range of about 1 to 5 amino acids. The variation allowed may be experimentally determined by producing the peptide synthetically while systematically making insertions, deletions, or substitutions of nucleotides in the sequence using recombinant DNA techniques.

30 A “signal sequence“ or “leader sequence“ can be used, when desired, to direct the polypeptide through a membrane of a cell. Such a sequence may be naturally present on the polypeptides of the present invention or provided from heterologous sources by recombinant DNA techniques.

An "oligopeptide" is a short stretch of amino acid residues and may be expressed from an oligonucleotide. Oligopeptides comprise a stretch of amino acid residues of at least 3, 5, 10 amino acids and at most 10, 15, 25 amino acids, typically of at least 9 to 13 amino acids, and of sufficient length to display biological and/or antigenic activity.

- 5 "Inhibitor" is any substance which retards or prevents a chemical or physiological reaction or response. Common inhibitors include but are not limited to antisense molecules, antibodies, and antagonists including small compound modulators of functional ion channel activity.

"Standard expression" is a quantitative or qualitative measurement for comparison. It is based on a statistically appropriate number of normal samples and is created to use as a basis of comparison
10 when performing diagnostic assays, running clinical trials, or following patient treatment profiles.

"Animal" as used herein may be defined to include human, domestic (e.g., cats, dogs, etc.), agricultural (e.g., cows, horses, sheep, etc.) or test species (e.g., mouse, rat, rabbit, etc.).

A "KCNK1 polynucleotide", within the meaning of the invention, shall be understood as being a nucleic acid molecule selected from a group consisting of

- 15 (i) nucleic acid molecules encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 2,
(ii) nucleic acid molecules comprising the sequence of SEQ ID NO: 1,
(iii) nucleic acid molecules having the sequence of SEQ ID NO: 1,
(iv) nucleic acid molecules the complementary strand of which hybridizes under stringent
20 conditions to a nucleic acid molecule of (i), (ii), or (iii); and
(v) nucleic acid molecules the sequence of which differs from the sequence of a nucleic acid molecule of (iii) due to the degeneracy of the genetic code;

wherein the polypeptide encoded by said nucleic acid molecule has KCNK1 activity.

A "KCNK1 polypeptide", within the meaning of the invention, shall be understood as being a
25 polypeptide selected from a group consisting of

- (i) polypeptides having the sequence of SEQ ID NO: 2,
(ii) polypeptides comprising the sequence of SEQ ID NO: 2,
(iii) polypeptides encoded by KCNK1 polynucleotides; and
(iv) polypeptides which show at least 99%, 98%, 95%, 90%, or 80% homology with a
30 polypeptide of (i), (ii), or (iii);

wherein said polypeptide has KCNK1 activity.

The nucleotide sequences encoding a KCNK1 (or their complement) have numerous applications in techniques known to those skilled in the art of molecular biology. These techniques include use as hybridization probes, use in the construction of oligonucleotides for PCR, use for chromosome and gene mapping, use in the recombinant production of KCNK1, and use in generation of antisense DNA or RNA, their chemical analogs and the like. Uses of nucleotides encoding a KCNK1 disclosed herein are exemplary of known techniques and are not intended to limit their use in any technique known to a person of ordinary skill in the art. Furthermore, the nucleotide sequences disclosed herein may be used in molecular biology techniques that have not yet been developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, e.g., the triplet genetic code, specific base pair interactions, etc.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of KCNK1 - encoding nucleotide sequences may be produced. Some of these will only bear minimal homology to the nucleotide sequence of the known and naturally occurring KCNK1. The invention has specifically contemplated each and every possible variation of nucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the nucleotide sequence of naturally occurring KCNK1, and all such variations are to be considered as being specifically disclosed.

Although the nucleotide sequences which encode a KCNK1, its derivatives or its variants are preferably capable of hybridizing to the nucleotide sequence of the naturally occurring KCNK1 polynucleotide under stringent conditions, it may be advantageous to produce nucleotide sequences encoding KCNK1 polypeptides or its derivatives possessing a substantially different codon usage. Codons can be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic expression host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding a KCNK1 polypeptide and/or its derivatives without altering the encoded amino acid sequence include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

Nucleotide sequences encoding a KCNK1 polypeptide may be joined to a variety of other nucleotide sequences by means of well established recombinant DNA techniques. Useful nucleotide sequences for joining to KCNK1 polynucleotides include an assortment of cloning vectors such as plasmids, cosmids, lambda phage derivatives, phagemids, and the like. Vectors of interest include expression vectors, replication vectors, probe generation vectors, sequencing

vectors, etc. In general, vectors of interest may contain an origin of replication functional in at least one organism, convenient restriction endonuclease sensitive sites, and selectable markers for one or more host cell systems.

Another aspect of the subject invention is to provide for KCNK1-specific hybridization probes
5 capable of hybridizing with naturally occurring nucleotide sequences encoding KCNK1. Such probes may also be used for the detection of similar ion channel encoding sequences and should preferably show at least 40% nucleotide identity to KCNK1 polynucleotides. The hybridization probes of the subject invention may be derived from the nucleotide sequence presented as SEQ ID NO: 1 or from genomic sequences including promoter, enhancers or introns of the native gene.
10 Hybridization probes may be labelled by a variety of reporter molecules using techniques well known in the art.

It will be recognized that many deletional or mutational analogs of KCNK1 polynucleotides will be effective hybridization probes for KCNK1 polynucleotides. Accordingly, the invention relates to nucleic acid sequences that hybridize with such KCNK1 encoding nucleic acid sequences under
15 stringent conditions.

“Stringent conditions” refers to conditions that allow for the hybridization of substantially related nucleic acid sequences. For instance, such conditions will generally allow hybridization of sequence with at least about 85% sequence identity, preferably with at least about 90% sequence identity, more preferably with at least about 95% sequence identity. Hybridization conditions and
20 probes can be adjusted in well-characterized ways to achieve selective hybridization of human-derived probes. Stringent conditions, within the meaning of the invention are 65°C in a buffer containing 1 mM EDTA, 0.5 M NaHPO₄ (pH 7.2), 7 % (w/v) SDS.

Nucleic acid molecules that will hybridize to KCNK1 polynucleotides under stringent conditions can be identified functionally. Without limitation, examples of the uses for hybridization probes
25 include: histochemical uses such as identifying tissues that express KCNK1; measuring mRNA levels, for instance to identify a sample's tissue type or to identify cells that express abnormal levels of KCNK1; and detecting polymorphisms of KCNK1.

PCR provides additional uses for oligonucleotides based upon the nucleotide sequence which encodes KCNK1. Such probes used in PCR may be of recombinant origin, chemically
30 synthesized, or a mixture of both. Oligonucleotides may comprise discrete nucleotide sequences employed under optimized conditions for identification of KCNK1 in specific tissues or diagnostic use. The same two oligomers, a nested set of oligonucleotides, or even a degenerate pool of

oligonucleotides may be employed under less stringent conditions for identification of closely related DNAs or RNAs.

Rules for designing polymerase chain reaction (PCR) primers are now established, as reviewed by PCR Protocols. Degenerate primers, i.e., preparations of primers that are heterogeneous at given
5 sequence locations, can be designed to amplify nucleic acid sequences that are highly homologous to, but not identical with KCNK1. Strategies are now available that allow for only one of the primers to be required to specifically hybridize with a known sequence. For example, appropriate nucleic acid primers can be ligated to the nucleic acid sought to be amplified to provide the hybridization partner for one of the primers. In this way, only one of the primers need be based on
10 the sequence of the nucleic acid sought to be amplified.

PCR methods for amplifying nucleic acid will utilize at least two primers. One of these primers will be capable of hybridizing to a first strand of the nucleic acid to be amplified and of priming enzyme-driven nucleic acid synthesis in a first direction. The other will be capable of hybridizing the reciprocal sequence of the first strand (if the sequence to be amplified is single stranded, this
15 sequence will initially be hypothetical, but will be synthesized in the first amplification cycle) and of priming nucleic acid synthesis from that strand in the direction opposite the first direction and towards the site of hybridization for the first primer. Conditions for conducting such amplifications, particularly under preferred stringent hybridization conditions, are well known.

Other means of producing specific hybridization probes for KCNK1 include the cloning of nucleic
20 acid sequences encoding KCNK1 or KCNK1 derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerase as T7 or SP6 RNA polymerase and the appropriate reporter molecules.

It is possible to produce a DNA sequence, or portions thereof, entirely by synthetic chemistry.
25 After synthesis, the nucleic acid sequence can be inserted into any of the many available DNA vectors and their respective host cells using techniques which are well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into the nucleotide sequence. Alternately, a portion of sequence in which a mutation is desired can be synthesized and recombined with longer portion of an existing genomic or recombinant sequence.

30 KCNK1 polynucleotides may be used to produce a purified oligo-or polypeptide using well known methods of recombinant DNA technology. The oligopeptide may be expressed in a variety of host cells, either prokaryotic or eukaryotic. Host cells may be from the same species from which the nucleotide sequence was derived or from a different species. Advantages of producing an

oligonucleotide by recombinant DNA technology include obtaining adequate amounts of the protein for purification and the availability of simplified purification procedures.

Quantitative determinations of nucleic acids

5 An important step in the molecular genetic analysis of human disease is often the enumeration of the copy number of a nucleic acid or the relative expression of a gene in particular tissues.

Several different approaches are currently available to make quantitative determinations of nucleic acids. Chromosome-based techniques, such as comparative genomic hybridization (CGH) and fluorescent in situ hybridization (FISH) facilitate efforts to cytogenetically localize genomic regions that are altered in tumor cells. Regions of genomic alteration can be narrowed further using loss of heterozygosity analysis (LOH), in which disease DNA is analyzed and compared with 10 normal DNA for the loss of a heterozygous polymorphic marker. The first experiments used restriction fragment length polymorphisms (RFLPs) [Johnson, (1989)], or hypervariable minisatellite DNA [Barnes, 2000]. In recent years LOH has been performed primarily using PCR amplification of microsatellite markers and electrophoresis of the radio labelled [Jeffreys, (1985)] 15 or fluorescently labelled PCR products [Weber, (1990)] and compared between paired normal and disease DNAs.

A number of other methods have also been developed to quantify nucleic acids [Gergen, (1992)]. More recently, PCR and RT-PCR methods have been developed which are capable of measuring the amount of a nucleic acid in a sample. One approach, for example, measures PCR product 20 quantity in the log phase of the reaction before the formation of reaction products plateaus [Thomas, (1980)].

A gene sequence contained in all samples at relatively constant quantity is typically utilized for sample amplification efficiency normalization. This approach, however, suffers from several drawbacks. The method requires that each sample has equal input amounts of the nucleic acid and 25 that the amplification efficiency between samples is identical until the time of analysis. Furthermore, it is difficult using the conventional methods of PCR quantitation such as gel electrophoresis or plate capture hybridization to determine that all samples are in fact analyzed during the log phase of the reaction as required by the method.

Another method called quantitative competitive (QC)-PCR, as the name implies, relies on the 30 inclusion of an internal control competitor in each reaction [Piatak, (1993), BioTechniques]. The efficiency of each reaction is normalized to the internal competitor. A known amount of internal competitor is typically added to each sample. The unknown target PCR product is compared with

the known competitor PCR product to obtain relative quantitation. A difficulty with this general approach lies in developing an internal control that amplifies with the same efficiency than the target molecule.

5' Fluorogenic Nuclease Assays

- 5 Fluorogenic nuclease assays are a real time quantitation method that uses a probe to monitor formation of amplification product. The basis for this method of monitoring the formation of amplification product is to measure continuously PCR product accumulation using a dual-labelled fluorogenic oligonucleotide probe, an approach frequently referred to in the literature simply as the "TaqMan method" [Piatak, (1993), Science; Heid, (1996); Gibson, (1996); Holland. (1991)].
- 10 The probe used in such assays is typically a short (about 20-25 bases) oligonucleotide that is labeled with two different fluorescent dyes. The 5' terminus of the probe is attached to a reporter dye and the 3' terminus is attached to a quenching dye, although the dyes could be attached at other locations on the probe as well. The probe is designed to have at least substantial sequence complementarity with the probe binding site. Upstream and downstream PCR primers which bind
- 15 to flanking regions of the locus are added to the reaction mixture. When the probe is intact, energy transfer between the two fluorophors occurs and the quencher quenches emission from the reporter. During the extension phase of PCR, the probe is cleaved by the 5' nuclease activity of a nucleic acid polymerase such as Taq polymerase, thereby releasing the reporter from the oligonucleotide-quencher and resulting in an increase of reporter emission intensity which can be
- 20 measured by an appropriate detector.

One detector which is specifically adapted for measuring fluorescence emissions such as those created during a fluorogenic assay is the ABI 7700 or 4700 HT manufactured by Applied Biosystems, Inc. in Foster City, Calif. The ABI 7700 uses fiber optics connected with each well in a 96-or 384 well PCR tube arrangement. The instrument includes a laser for exciting the labels

25 and is capable of measuring the fluorescence spectra intensity from each tube with continuous monitoring during PCR amplification. Each tube is re-examined every 8.5 seconds.

Computer software provided with the instrument is capable of recording the fluorescence intensity of reporter and quencher over the course of the amplification. The recorded values will then be used to calculate the increase in normalized reporter emission intensity on a continuous basis. The

30 increase in emission intensity is plotted versus time, i.e., the number of amplification cycles, to produce a continuous measure of amplification. To quantify the locus in each amplification reaction, the amplification plot is examined at a point during the log phase of product accumulation. This is accomplished by assigning a fluorescence threshold intensity above background

and determining the point at which each amplification plot crosses the threshold (defined as the threshold cycle number or Ct). Differences in threshold cycle number are used to quantify the relative amount of PCR target contained within each tube. Assuming that each reaction functions at 100% PCR efficiency, a difference of one Ct represents a two-fold difference in the amount of starting template. The fluorescence value can be used in conjunction with a standard curve to determine the amount of amplification product present.

Non-Probe-Based Detection Methods

A variety of options are available for measuring the amplification products as they are formed. One method utilizes labels, such as dyes, which only bind to double stranded DNA. In this type of approach, amplification product (which is double stranded) binds dye molecules in solution to form a complex. With the appropriate dyes, it is possible to distinguish between dye molecules free in solution and dye molecules bound to amplification product. For example, certain dyes fluoresce only when bound to amplification product. Examples of dyes which can be used in methods of this general type include, but are not limited to, Syber Green.TM. and Pico Green from Molecular Probes, Inc. of Eugene, Oreg., ethidium bromide, propidium iodide, chromomycin, acridine orange, Hoechst 33258, Toto-1, Yoyo-1, DAPI (4',6-diamidino-2-phenylindole hydrochloride).

Another real time detection technique measures alteration in energy fluorescence energy transfer between fluorophors conjugated with PCR primers [Livak, (1995)].

Probe-Based Detection Methods

These detection methods involve some alteration to the structure or conformation of a probe hybridized to the locus between the amplification primer pair. In some instances, the alteration is caused by the template-dependent extension catalyzed by a nucleic acid polymerase during the amplification process. The alteration generates a detectable signal which is an indirect measure of the amount of amplification product formed.

For example, some methods involve the degradation or digestion of the probe during the extension reaction. These methods are a consequence of the 5'-3' nuclease activity associated with some nucleic acid polymerases. Polymerases having this activity cleave mononucleotides or small oligonucleotides from an oligonucleotide probe annealed to its complementary sequence located within the locus.

The 3' end of the upstream primer provides the initial binding site for the nucleic acid polymerase. As the polymerase catalyzes extension of the upstream primer and encounters the bound probe, the

nucleic acid polymerase displaces a portion of the 5' end of the probe and through its nuclease activity cleaves mononucleotides or oligonucleotides from the probe.

The upstream primer and the probe can be designed such that they anneal to the complementary strand in close proximity to one another. In fact, the 3' end of the upstream primer and the 5' end of the probe may abut one another. In this situation, extension of the upstream primer is not necessary in order for the nucleic acid polymerase to begin cleaving the probe. In the case in which intervening nucleotides separate the upstream primer and the probe, extension of the primer is necessary before the nucleic acid polymerase encounters the 5' end of the probe. Once contact occurs and polymerization continues, the 5'-3' exonuclease activity of the nucleic acid polymerase begins cleaving mononucleotides or oligonucleotides from the 5' end of the probe. Digestion of the probe continues until the remaining portion of the probe dissociates from the complementary strand.

In solution, the two end sections can hybridize with each other to form a hairpin loop. In this conformation, the reporter and quencher dye are in sufficiently close proximity that fluorescence from the reporter dye is effectively quenched by the quencher dye. Hybridized probe, in contrast, results in a linearized conformation in which the extent of quenching is decreased. Thus, by monitoring emission changes for the two dyes, it is possible to indirectly monitor the formation of amplification product.

Probes

The labeled probe is selected so that its sequence is substantially complementary to a segment of the test locus or a reference locus. As indicated above, the nucleic acid site to which the probe binds should be located between the primer binding sites for the upstream and downstream amplification primers.

Primers

The primers used in the amplification are selected so as to be capable of hybridizing to sequences at flanking regions of the locus being amplified. The primers are chosen to have at least substantial complementarity with the different strands of the nucleic acid being amplified. When a probe is utilized to detect the formation of amplification products, the primers are selected in such that they flank the probe, i.e. are located upstream and downstream of the probe.

The primer must have sufficient length so that it is capable of priming the synthesis of extension products in the presence of an agent for polymerization. The length and composition of the primer depends on many parameters, including, for example, the temperature at which the annealing

reaction is conducted, proximity of the probe binding site to that of the primer, relative concentrations of the primer and probe and the particular nucleic acid composition of the probe. Typically the primer includes 15-30 nucleotides. However, the length of the primer may be more or less depending on the complexity of the primer binding site and the factors listed above.

5 *Labels for Probes and Primers*

The labels used for labeling the probes or primers of the current invention and which can provide the signal corresponding to the quantity of amplification product can take a variety of forms. As indicated above with regard to the 5' fluorogenic nuclease method, a fluorescent signal is one signal which can be measured. However, measurements may also be made, for example, by
10 monitoring radioactivity, colorimetry, absorption, magnetic parameters, or enzymatic activity. Thus, labels which can be employed include, but are not limited to, fluorophors, chromophores, radioactive isotopes, electron dense reagents, enzymes, and ligands having specific binding partners (e.g., biotin-avidin).

Monitoring changes in fluorescence is a particularly useful way to monitor the accumulation of
15 amplification products. A number of labels useful for attachment to probes or primers are commercially available including fluorescein and various fluorescein derivatives such as FAM, HEX, TET and JOE (all which are available from Applied Biosystems, Foster City, Calif.); lucifer yellow, and coumarin derivatives.

Labels may be attached to the probe or primer using a variety of techniques and can be attached at
20 the 5' end, and/or the 3' end and/or at an internal nucleotide. The label can also be attached to spacer arms of various sizes which are attached to the probe or primer. These spacer arms are useful for obtaining a desired distance between multiple labels attached to the probe or primer.

In some instances, a single label may be utilized; whereas, in other instances, such as with the 5' fluorogenic nuclease assays for example, two or more labels are attached to the probe. In cases
25 wherein the probe includes multiple labels, it is generally advisable to maintain spacing between the labels which is sufficient to permit separation of the labels during digestion of the probe through the 5'-3' nuclease activity of the nucleic acid polymerase.

Patients Exhibiting Symptoms of Disease

A number of diseases are associated with changes in the copy number of a certain gene. For
30 patients having symptoms of a disease, the real-time PCR method can be used to determine if the patient has copy number alterations which are known to be linked with diseases that are associated with the symptoms the patient has.

KCNK1 expression

KCNK1 fusion proteins

Fusion proteins are useful for generating antibodies against KCNK1 polypeptides and for use in various assay systems. For example, fusion proteins can be used to identify proteins which
5 interact with portions of KCNK1 polypeptides. Protein affinity chromatography or library-based assays for protein-protein interactions, such as the yeast two-hybrid or phage display systems, can be used for this purpose. Such methods are well known in the art and also can be used as drug screens.

A KCNK1 fusion protein comprises two polypeptide segments fused together by means of a
10 peptide bond. The first polypeptide segment can comprise at least 54, 75, 100, 125, 139, 150, 175, 200, 225, 250, or 275 contiguous amino acids of SEQ ID NO: 2 or of a biologically active variant, such as those described above. The first polypeptide segment also can comprise full-length KCNK1.

The second polypeptide segment can be a full-length protein or a protein fragment. Proteins
15 commonly used in fusion protein construction include, but are not limited to β galactosidase, β -glucuronidase, green fluorescent protein (GFP), autofluorescent proteins, including blue fluorescent protein (BFP), glutathione-S-transferase (GST), luciferase, horseradish peroxidase (HRP), and chloramphenicol acetyltransferase (CAT). Additionally, epitope tags are used in fusion protein constructions, including histidine (His) tags, FLAG tags, influenza hemagglutinin (HA)
20 tags, Myc tags, VSV-G tags, and thioredoxin (Trx) tags. Other fusion constructions can include maltose binding protein (MBP), S-tag, Lex a DNA binding domain (DBD) fusions, GAL4 DNA binding domain fusions, or herpes simplex virus (HSV) BP16 protein fusions. A fusion protein also can be engineered to contain a cleavage site located adjacent to the KCNK1.

Preparation of Polynucleotides

25 A naturally occurring KCNK1 polynucleotide can be isolated free of other cellular components such as membrane components, proteins, and lipids. Polynucleotides can be made by a cell and isolated using standard nucleic acid purification techniques, or synthesized using an amplification technique, such as the polymerase chain reaction (PCR), or by using an automatic synthesizer. Methods for isolating polynucleotides are routine and are known in the art. Any such technique
30 for obtaining a polynucleotide can be used to obtain isolated KCNK1 polynucleotides. For example, restriction enzymes and probes can be used to isolate polynucleotide fragments which

comprise KCNK1 nucleotide sequences. Isolated polynucleotides are in preparations which are free or at least 70, 80, or 90% free of other molecules.

KCNK1 cDNA molecules can be made with standard molecular biology techniques, using KCNK1 mRNA as a template. KCNK1 cDNA molecules can thereafter be replicated using molecular
5 biology techniques known in the art. An amplification technique, such as PCR, can be used to obtain additional copies of polynucleotides of the invention, using either human genomic DNA or cDNA as a template.

Alternatively, synthetic chemistry techniques can be used to synthesize KCNK1 polynucleotides. The degeneracy of the genetic code allows alternate nucleotide sequences to be synthesized which
10 will encode KCNK1 having, for example, an amino acid sequence shown in SEQ ID NO: 2 or a biologically active variant thereof.

Extending Polynucleotides

Various PCR-based methods can be used to extend nucleic acid sequences encoding human KCNK1, for example to detect upstream sequences of KCNK1 gene such as promoters and
15 regulatory elements. For example, restriction-site PCR uses universal primers to retrieve unknown sequence adjacent to a known locus. Genomic DNA is first amplified in the presence of a primer to a linker sequence and a primer specific to the known region. The amplified sequences are then subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA
20 polymerase and sequenced using reverse transcriptase.

Inverse PCR also can be used to amplify or extend sequences using divergent primers based on a known region. Primers can be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences Inc., Plymouth, Minn.), to be 22-30
25 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68-72°C. The method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

Another method which can be used is capture PCR, which involves PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial chromosome DNA. In this
30 method, multiple restriction enzyme digestions and ligations also can be used to place an engineered double-stranded sequence into an unknown fragment of the DNA molecule before performing PCR.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. Randomly-primed libraries are preferable, in that they will contain more sequences which contain the 5' regions of genes. Use of a randomly primed library may be especially preferable for situations in which an oligo d(T) library does not yield a full-length
5 cDNA. Genomic libraries can be useful for extension of sequence into 5' non-transcribed regulatory regions.

Commercially available capillary electrophoresis systems can be used to analyze the size or confirm the nucleotide sequence of PCR or sequencing products. For example, capillary sequencing can employ flowable polymers for electrophoretic separation, four different fluorescent
10 dyes (one for each nucleotide) which are laser activated, and detection of the emitted wavelengths by a charge coupled device camera. Output/light intensity can be converted to electrical signal using appropriate equipment and software (e.g., GENOTYPER and Sequence NAVIGATOR, Perkin Elmer), and the entire process from loading of samples to computer analysis and electronic data display can be computer controlled. Capillary electrophoresis is especially preferable for the
15 sequencing of small pieces of DNA which might be present in limited amounts in a particular sample.

Obtaining Polypeptides

KCNK1 can be obtained, for example, by purification from human cells, by expression of KCNK1 polynucleotides, or by direct chemical synthesis.

20 *Protein Purification*

KCNK1 can be purified from any human cell which expresses the channel, including those which have been transfected with expression constructs which express KCNK1. A purified KCNK1 is separated from other compounds which normally associate with KCNK1 in the cell, such as certain proteins, carbohydrates, or lipids, using methods well-known in the art. Such methods
25 include, but are not limited to, size exclusion chromatography, ammonium sulfate fractionation, ion exchange chromatography, affinity chromatography, and preparative gel electrophoresis.

Expression of KCNK1 Polynucleotides

To express KCNK1, KCNK1 polynucleotides can be inserted into an expression vector which contains the necessary elements for the transcription and translation of the inserted coding
30 sequence. Methods which are well known to those skilled in the art can be used to construct expression vectors containing sequences encoding KCNK1 and appropriate transcriptional and

translational control elements. These methods include *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination.

A variety of expression vector/host systems can be utilized to contain and express sequences encoding KCNK1. These include, but are not limited to, microorganisms, such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors, insect cell systems infected with virus expression vectors (*e.g.*, baculovirus), plant cell systems transformed with virus expression vectors (*e.g.*, cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (*e.g.*, Ti or pBR322 plasmids), or animal cell systems.

10 The control elements or regulatory sequences are those non-translated regions of the vector - enhancers, promoters, 5' and 3' untranslated regions -- which interact with host cellular proteins to carry out transcription and translation. Such elements can vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, can be used. For example, 15 when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the BLUESCRIPT phagemid (Stratagene, LaJolla, Calif.) or pSPORT1 plasmid (Life Technologies) and the like can be used. The baculovirus polyhedrin promoter can be used in insect cells. Promoters or enhancers derived from the genomes of plant cells (*e.g.*, heat shock, RUBISCO, and storage protein genes) or from plant viruses (*e.g.*, viral promoters or leader sequences) can be 20 cloned into the vector. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are preferable. If it is necessary to generate a cell line that contains multiple copies of a nucleotide sequence encoding KCNK1, vectors based on SV40 or EBV can be used with an appropriate selectable marker.

Bacterial and Yeast Expression Systems

25 In bacterial systems, a number of expression vectors can be selected. For example, when a large quantity of KCNK1 is needed for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified can be used. Such vectors include, but are not limited to, multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene). In a BLUESCRIPT vector, a sequence encoding KCNK1 can be ligated into the 30 vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced. pIN vectors or pGEX vectors (Promega, Madison, Wis.) also can be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the

presence of free glutathione. Proteins made in such systems can be designed to include heparin, thrombin, or factor Xa protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

Plant and Insect Expression Systems

5 If plant expression vectors are used, the expression of sequences encoding KCNK1 can be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV can be used alone or in combination with the omega leader sequence from TMV. Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters can be used. These constructs can be introduced into plant cells by direct DNA
10 transformation or by pathogen-mediated transfection.

An insect system also can be used to express KCNK1. For example, in one such system *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. Sequences encoding KCNK1 can be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under
15 control of the polyhedrin promoter. Successful insertion of KCNK1 will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses can then be used to infect *S. frugiperda* cells or *Trichoplusia* larvae in which KCNK1 can be expressed.

Mammalian Expression Systems

20 A number of viral-based expression systems can be used to express KCNK1 in mammalian host cells. For example, if an adenovirus is used as an expression vector, sequences encoding KCNK1 can be ligated into an adenovirus transcription/translation complex comprising the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome can be used to obtain a viable virus which is capable of expressing KCNK1 in infected host cells
25 [Engelhard, 1994]. If desired, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, can be used to increase expression in mammalian host cells.

Human artificial chromosomes (HACs) also can be used to deliver larger fragments of DNA than can be contained and expressed in a plasmid. HACs of 6M to 10M are constructed and delivered to cells via conventional delivery methods (e.g., liposomes, polycationic amino polymers, or
30 vesicles). Specific initiation signals also can be used to achieve more efficient translation of sequences encoding KCNK1. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding KCNK1, its initiation codon, and upstream

sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals (including the ATG initiation codon) should be provided. The initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons can be of various origins, both natural and synthetic.

Host Cells

A host cell strain can be chosen for its ability to modulate the expression of the inserted sequences, to process the expressed KCNK1 in the desired fashion, or to improve the readout window for the desired KCNK1. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the polypeptide also can be used to facilitate correct insertion, folding and/or function. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (*e.g.*, CHO, HeLa, MDCK, HEK293, and WI38), are available from the American Type Culture Collection (ATCC; 10801 University Boulevard, Manassas, VA 20110-2209) and can be chosen to ensure the correct modification and processing of the foreign protein.

Stable expression is preferred for long-term, high-yield production of recombinant proteins. For example, cell lines which stably express KCNK1 can be transformed using expression vectors which can contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells can be allowed to grow for 1-2 days in an enriched medium before they are switched to a selective medium. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced KCNK1 sequences. Resistant clones of stably transformed cells can be proliferated using tissue culture techniques appropriate to the cell type. Any number of selection systems can be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase [Logan, (1984)] and adenine phosphoribosyltransferase [Wigler, (1977)] genes which can be employed in *tk* or *aprt* cells, respectively. Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate [Lowy, (1980)], *npt* confers resistance to the aminoglycosides, neomycin and G-418 [Wigler, (1980)], and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively [Colbere-Garapin, 1981]. Additional selectable genes have been described. For example, *trpB* allows cells to utilize indole in place of tryptophan, or *hisD*, which

allows cells to utilize histinol in place of histidine. Visible markers such as anthocyanins, β -glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, can be used to identify transformants and to quantify the amount of transient or stable protein expression attributable to a specific vector system

5 *Detecting Polypeptide Expression*

Although the presence of marker gene expression suggests that a KCNK1 polynucleotide is also present, its presence and expression may need to be confirmed. For example, if a sequence encoding KCNK1 is inserted within a marker gene sequence, transformed cells containing sequences which encode KCNK1 can be identified by the absence of marker gene function.
10 Alternatively, a marker gene can be placed in tandem with a sequence encoding KCNK1 under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of KCNK1 polynucleotide.

Alternatively, host cells which contain a KCNK1 polynucleotide and which express KCNK1 can be identified by a variety of procedures known to those of skill in the art. These procedures
15 include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassay or immunoassay techniques which include membrane, solution, or chip-based technologies for the detection and/or quantification of nucleic acid or protein. For example, the presence of a polynucleotide sequence encoding KCNK1 can be detected by DNA-DNA or DNA-RNA hybridization or amplification using probes or fragments or fragments of polynucleotides encoding KCNK1.
20 Nucleic acid amplification-based assays involve the use of oligonucleotides selected from sequences encoding KCNK1 to detect transformants which contain a KCNK1 polynucleotide.

A variety of protocols for detecting and measuring the expression of KCNK1, using either polyclonal or monoclonal antibodies specific for the polypeptide, are known in the art. Examples
25 include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay using monoclonal antibodies reactive to two non-interfering epitopes on KCNK1 can be used, or a competitive binding assay can be employed.

A wide variety of labels and conjugation techniques are known by those skilled in the art and can be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization
30 or PCR probes for detecting sequences related to polynucleotides encoding KCNK1 include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, sequences encoding KCNK1 can be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and can be used to

synthesize RNA probes *in vitro* by addition of labeled nucleotides and an appropriate RNA polymerase such as T7, T3, or SP6. These procedures can be conducted using a variety of commercially available kits (Amersham Pharmacia Biotech, Promega, and US Biochemical). Suitable reporter molecules or labels which can be used for ease of detection include
5 radionuclides, enzymes, and fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Expression and Purification of Polypeptides

Host cells transformed with KCNK1 polynucleotides can be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The polypeptide produced by a
10 transformed cell can be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing KCNK1 polynucleotides can be designed to contain signal sequences which direct secretion of soluble KCNK1 through a prokaryotic or eukaryotic cell membrane or which direct the membrane insertion of membrane-bound KCNK1.

15 As discussed above, other constructions can be used to join a sequence encoding KCNK1 to a nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain
20 utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Wash.). Inclusion of cleavable linker sequences such as those specific for Factor XA or enterokinase (Invitrogen, San Diego, CA) between the purification domain and KCNK1 also can be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing KCNK1 and 6 histidine residues preceding a thioredoxin or an enterokinase cleavage
25 site. The histidine residues facilitate purification by IMAC (immobilized metal ion affinity chromatography) Maddox, (1983)], while the enterokinase cleavage site provides a means for purifying KCNK1 from the fusion protein [Porath, (1992)].

Chemical Synthesis

Sequences encoding KCNK1 can be synthesized, in whole or in part, using chemical methods well
30 known in the art. Alternatively, KCNK1 itself can be produced using chemical methods to synthesize its amino acid sequence, such as by direct peptide synthesis using solid-phase techniques. Protein synthesis can either be performed using manual techniques or by automation. Automated synthesis can be achieved, for example, using Applied Biosystems 431A Peptide

Synthesizer (Perkin Elmer). Optionally, fragments of KCNK1 can be separately synthesized and combined using chemical methods to produce a full-length molecule.

The newly synthesized peptide can be substantially purified by preparative high performance liquid chromatography. The composition of a synthetic KCNK1 can be confirmed by amino acid
5 analysis or sequencing. Additionally, any portion of the amino acid sequence of KCNK1 can be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins to produce a variant polypeptide or a fusion protein.

Production of Altered Polypeptides

As will be understood by those of skill in the art, it may be advantageous to produce KCNK1
10 polynucleotides possessing non-naturally occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce an RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

The nucleotide sequences referred to herein can be engineered using methods generally known in
15 the art to alter KCNK1 polynucleotides for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the polypeptide or mRNA product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides can be used to engineer the nucleotide sequences. For example, site-directed mutagenesis can be used to insert new restriction sites, alter glycosylation patterns,
20 change codon preference, produce splice variants, introduce mutations, and so forth.

Antibodies

Any type of antibody known in the art can be generated to bind specifically to an epitope of KCNK1.

"Antibody" as used herein includes intact immunoglobulin molecules, as well as fragments
25 thereof, such as Fab, F(ab')₂, and Fv, which are capable of binding an epitope of KCNK1. Typically, at least 6, 8, 10, or 12 contiguous amino acids are required to form an epitope. However, epitopes which involve non-contiguous amino acids may require more, e.g., at least 15, 25, or 50 amino acid. An antibody which specifically binds to an epitope of KCNK1 can be used therapeutically, as well as in immunochemical assays, such as Western blots, ELISAs, radioimmuno-
30 assays, immunohistochemical assays, immunoprecipitations, or other immunochemical assays known in the art. Various immunoassays can be used to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays are well

known in the art. Such immunoassays typically involve the measurement of complex formation between an immunogen and an antibody which specifically binds to the KCNK1 immunogen.

Typically, an antibody which specifically binds to KCNK1 provides a detection signal at least 5-, 10-, or 20-fold higher than a detection signal provided with other proteins when used in an immunochemical assay. Preferably, antibodies which specifically bind to KCNK1 do not detect other proteins in immunochemical assays and can immunoprecipitate KCNK1 from solution.

KCNK1 can be used to immunize a mammal, such as a mouse, rat, rabbit, guinea pig, monkey, or human, to produce polyclonal antibodies. If desired, KCNK1 can be conjugated to a carrier protein, such as bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin. Depending on the host species, various adjuvants can be used to increase the immunological response. Such adjuvants include, but are not limited to, Freund's adjuvant, mineral gels (e.g., aluminum hydroxide), and surface active substances (e.g., lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinitrophenol). Among adjuvants used in humans, BCG (*bacilli Calmette-Guerin*) and *Corynebacterium parvum* are especially useful.

Monoclonal antibodies which specifically bind to KCNK1 can be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These techniques include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique [Roberge, (1995)].

In addition, techniques developed for the production of "chimeric antibodies", the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. Monoclonal and other antibodies also can be "humanized" to prevent a patient from mounting an immune response against the antibody when it is used therapeutically. Such antibodies may be sufficiently similar in sequence to human antibodies to be used directly in therapy or may require alteration of a few key residues. Sequence differences between rodent antibodies and human sequences can be minimized by replacing residues which differ from those in the human sequences by site directed mutagenesis of individual residues or by grating of entire complementarity determining regions. Antibodies which specifically bind to KCNK1 can contain antigen binding sites which are either partially or fully humanized, as disclosed in U.S. 5,565,332.

Alternatively, techniques described for the production of single chain antibodies can be adapted using methods known in the art to produce single chain antibodies which specifically bind to KCNK1. Antibodies with related specificity, but of distinct idiotypic composition, can be generated by chain shuffling from random combinatorial immunoglobulin libraries. Single-chain

antibodies also can be constructed using a DNA amplification method, such as PCR, using hybridoma cDNA as a template. Single-chain antibodies can be mono- or bispecific, and can be bivalent or tetravalent. Construction of tetravalent, bispecific single-chain antibodies is taught. A nucleotide sequence encoding a single-chain antibody can be constructed using manual or automated nucleotide synthesis, cloned into an expression construct using standard recombinant DNA methods, and introduced into a cell to express the coding sequence, as described below. Alternatively, single-chain antibodies can be produced directly using, for example, filamentous phage technology.

Antibodies which specifically bind to KCNK1 also can be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents. Other types of antibodies can be constructed and used therapeutically in methods of the invention. For example, chimeric antibodies can be constructed as disclosed in WO 93/03151. Binding proteins which are derived from immunoglobulins and which are multivalent and multispecific, such as the "diabodies" described in WO 94/13804, also can be prepared.

Antibodies according to the invention can be purified by methods well known in the art. For example, antibodies can be affinity purified by passage over a column to which KCNK1 is bound. The bound antibodies can then be eluted from the column using a buffer with a high salt concentration.

20 *Antisense Oligonucleotides*

Antisense oligonucleotides are nucleotide sequences which are complementary to a specific DNA or RNA sequence. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form complexes and block either transcription or translation. Preferably, an antisense oligonucleotide is at least 11 nucleotides in length, but can be at least 12, 15, 20, 25, 30, 35, 40, 45, or 50 or more nucleotides long. Longer sequences also can be used. Antisense oligonucleotide molecules can be provided in a DNA construct and introduced into a cell as described above to decrease the level of KCNK1 gene products in the cell.

Antisense oligonucleotides can be deoxyribonucleotides, ribonucleotides, or a combination of both. Oligonucleotides can be synthesized manually or by an automated synthesizer, by covalently linking the 5' end of one nucleotide with the 3' end of another nucleotide with non-phosphodiester internucleotide linkages such as alkylphosphonates, phosphorothioates, phosphorodithioates, alkylphosphonothioates, alkylphosphonates, phosphoramidates, phosphate esters, carbamates, acetamidate, carboxymethyl esters, carbonates, and phosphate triesters.

Modifications of KCNK1 gene expression can be obtained by designing antisense oligonucleotides which will form duplexes to the control, 5', or regulatory regions of the KCNK1 gene. Oligonucleotides derived from the transcription initiation site, *e.g.*, between positions -10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using "triple helix" base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or chaperons. Therapeutic advances using triplex DNA have been described in the literature [Nicholls, (1993)]. An antisense oligonucleotide also can be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

10 Precise complementarity is not required for successful complex formation between an antisense oligonucleotide and the complementary sequence of a KCNK1 polynucleotide. Antisense oligonucleotides which comprise, for example, 2, 3, 4, or 5 or more stretches of contiguous nucleotides which are precisely complementary to a KCNK1 polynucleotide, each separated by a stretch of contiguous nucleotides which are not complementary to adjacent KCNK1 nucleotides, can provide

15 sufficient targeting specificity for KCNK1 mRNA. Preferably, each stretch of complementary contiguous nucleotides is at least 4, 5, 6, 7, or 8 or more nucleotides in length. Non-complementary intervening sequences are preferably 1, 2, 3, or 4 nucleotides in length. One skilled in the art can easily use the calculated melting point of an antisense-sense pair to determine the degree of mismatching which will be tolerated between a particular antisense oligonucleotide and a

20 particular KCNK1 polynucleotide sequence. Antisense oligonucleotides can be modified without affecting their ability to hybridize to a KCNK1 polynucleotide. These modifications can be internal or at one or both ends of the antisense molecule. For example, internucleoside phosphate linkages can be modified by adding cholesteryl or diamine moieties with varying numbers of carbon residues between the amino groups and terminal ribose. Modified bases and/or sugars,

25 such as arabinose instead of ribose, or a 3', 5'-substituted oligonucleotide in which the 3' hydroxyl group or the 5' phosphate group are substituted, also can be employed in a modified antisense oligonucleotide. These modified oligonucleotides can be prepared by methods well known in the art.

Ribozymes

30 Ribozymes are RNA molecules with catalytic activity [Uhlmann, (1987)]. Ribozymes can be used to inhibit gene function by cleaving an RNA sequence, as is known in the art. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. Examples include engineered hammerhead motif ribozyme molecules that can specifically and efficiently catalyze endo-

nucleolytic cleavage of specific nucleotide sequences. The coding sequence of a KCNK1 polynucleotide can be used to generate ribozymes which will specifically bind to mRNA transcribed from a KCNK1 polynucleotide. Methods of designing and constructing ribozymes which can cleave other RNA molecules in trans in a highly sequence specific manner have been developed and described in the art. For example, the cleavage activity of ribozymes can be targeted to specific RNAs by engineering a discrete "hybridization" region into the ribozyme. The hybridization region contains a sequence complementary to the target RNA and thus specifically hybridizes with the target RNA.

Specific ribozyme cleavage sites within a KCNK1 RNA target can be identified by scanning the target molecule for ribozyme cleavage sites which include the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target RNA containing the cleavage site can be evaluated for secondary structural features which may render the target inoperable. Suitability of candidate KCNK1 RNA targets also can be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays. The nucleotide sequences shown in SEQ ID NO: 1 and its complement provide sources of suitable hybridization region sequences. Longer complementary sequences can be used to increase the affinity of the hybridization sequence for the target. The hybridizing and cleavage regions of the ribozyme can be integrally related such that upon hybridizing to the target RNA through the complementary regions, the catalytic region of the ribozyme can cleave the target.

Ribozymes can be introduced into cells as part of a DNA construct. Mechanical methods, such as microinjection, liposome-mediated transfection, electroporation, or calcium phosphate precipitation, can be used to introduce a ribozyme-containing DNA construct into cells in which it is desired to decrease KCNK1 expression. Alternatively, if it is desired that the cells stably retain the DNA construct, the construct can be supplied on a plasmid and maintained as a separate element or integrated into the genome of the cells, as is known in the art. A ribozyme-encoding DNA construct can include transcriptional regulatory elements, such as a promoter element, an enhancer or UAS element, and a transcriptional terminator signal, for controlling transcription of ribozymes in the cells (U.S. 5,641,673). Ribozymes also can be engineered to provide an additional level of regulation, so that destruction of mRNA occurs only when both a ribozyme and a target gene are induced in the cells.

Screening / Screening Assays

Regulators

Regulators as used herein, refer to compounds that affect the functional activity of a KCNK1 in vitro and/or in vivo. Regulators can be agonists and antagonists of a KCNK1 polypeptide and can be compounds that exert their effect on the KCNK1 activity via the expression, via post-translational modifications, by direct interaction with the channel protein or by other means. Agonists of KCNK1 are molecules which, when bound to KCNK1, increase or prolong the functional activity of KCNK1. Agonists of KCNK1 include proteins, nucleic acids, carbohydrates, small molecules, or any other molecule which activate KCNK1. Antagonists of KCNK1 are molecules which, when bound to KCNK1, decrease the amount or the duration of the functional activity of KCNK1. Antagonists include proteins, nucleic acids, carbohydrates, antibodies, small molecules, or any other molecule which decrease the activity of KCNK1.

The term "modulate", as it appears herein, refers to a change in the activity of KCNK1 polypeptide. For example, modulation may cause an increase or a decrease in functional activity, binding characteristics, or any other biological, functional, or immunological properties of KCNK1.

As used herein, the terms "specific binding" or "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, or an antagonist. The interaction is dependent upon the presence of a particular structure of the protein recognized by the binding molecule (i.e., the antigenic determinant or epitope). For example, if an antibody is specific for epitope "A" the presence of a polypeptide containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The invention provides methods (also referred to herein as "screening assays") for identifying compounds which can be used for the treatment of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases. The methods entail the identification of candidate or test compounds or agents (e.g., peptides, peptidomimetics, small molecules or other molecules) which bind to KCNK1 and/or have a stimulatory or inhibitory effect on the biological functional activity of KCNK1 or its expression and then determining which of these compounds have an effect on symptoms or diseases regarding cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in an *in vivo* assay.

Candidate or test compounds or agents which bind to KCNK1 and/or have a stimulatory or inhibitory effect on the functional activity or the expression of KCNK1 are identified either in assays that employ cells which express KCNK1 on the cell surface (cell-based assays) or in assays with isolated KCNK1 (cell-free assays). The various assays can employ a variety of variants of KCNK1 (e.g., full-length KCNK1, a biologically active fragment of KCNK1, or a fusion protein which includes all or a portion of KCNK1). Moreover, KCNK1 can be derived from any suitable mammalian species (e.g., human KCNK1, rat KCNK1 or murine KCNK1). The assay can be a binding assay entailing direct or indirect measurement of the binding of a test compound or a known KCNK1 ligand to KCNK1. The assay can also be an activity assay entailing direct or indirect measurement of the activity of KCNK1. The assay can also be an expression assay entailing direct or indirect measurement of the expression of KCNK1 mRNA or KCNK1 protein. The various screening assays are combined with an *in vivo* assay entailing measuring the effect of the test compound on the symptoms of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases.

In one embodiment, the invention provides assays for screening candidate or test compounds which bind to or modulate the functional activity of a membrane-bound (cell surface expressed) form of KCNK1. Such assays can employ full-length KCNK1, a biologically active fragment of KCNK1, or a fusion protein which includes all or a portion of KCNK1. As described in greater detail below, the test compound can be obtained by any suitable means, e.g., from conventional compound libraries. Determining the ability of the test compound to bind to a membrane-bound form of KCNK1 can be accomplished, for example, by coupling the test compound with a radioisotope or enzymatic label such that binding of the test compound to the KCNK1 -expressing cell can be measured by detecting the labeled compound in a complex. For example, the test compound can be labelled with ^{125}I , ^{35}S , ^{14}C , or ^3H , either directly or indirectly, and the radioisotope detected by direct counting of radioemmission or by scintillation counting. Alternatively, the test compound can be enzymatically labelled with, for example, horseradish peroxidase, alkaline phosphatase, or luciferase, and the enzymatic label detected by determination of conversion of an appropriate substrate to product.

In a competitive binding format, the assay comprises contacting KCNK1 expressing cell with a known compound which carries a detectable label and which binds to KCNK1 to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with the KCNK1 expressing cell, wherein determining the ability of the test compound to interact with the KCNK1 expressing cell comprises determining the ability of the test compound to preferentially bind the KCNK1 expressing cell as compared to the known compound.

In another embodiment, the assay is a cell-based assay comprising contacting a cell expressing a membrane-bound form of KCNK1 (e.g., full-length KCNK1, a biologically active fragment of KCNK1, or a fusion protein which includes all or a portion of KCNK1) expressed on the cell surface with a test compound and determining the ability of the test compound to modulate (e.g., stimulate or inhibit) the functional activity of the membrane-bound form of KCNK1. Determining the ability of the test compound to modulate the functional activity of the membrane-bound form of KCNK1 can be accomplished by any method suitable for measuring the functional activity of KCNK1, e.g., any method suitable for measuring the activity of any ion channel or other electrogenic target protein (described in greater detail below). The activity of an electrogenic target can be measured in a number of ways, not all of which are suitable for any given target protein. Among the measures of activity are: alteration in ion concentrations (e.g. Terstappen, 1999 *Analyt. Biochem.* 272, 149-155; Berman et al., 2000 *J Neurochem* 74(4), 1443-1451), measuring alterations in microphysiometric conditions (Hahnenberger et al., *Nature Biotech* 14, 880-884), measuring alterations in voltage or measuring alterations in currents (e.g. Gonzalez et al., 1999 4(9), 431-439; Willumsen et al., 2003 *Receptors and Channels* 9, 3-12).

The present invention also includes cell-free assays. Such assays involve contacting a form of KCNK1 (e.g., full-length KCNK1, a biologically active fragment of KCNK1, or a fusion protein comprising all or a portion of KCNK1) with a test compound and determining the ability of the test compound to bind to KCNK1. Binding of the test compound to KCNK1 can be determined either directly or indirectly as described above. In one embodiment, the assay includes contacting KCNK1 with a known compound which carries a detectable label and which binds KCNK1 to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with KCNK1, wherein determining the ability of the test compound to interact with KCNK1 comprises determining the ability of the test compound to preferentially bind to KCNK1 as compared to the known compound.

The cell-free assays of the present invention are amenable to use of either a membrane-bound form of KCNK1 or a soluble fragment thereof. In the case of cell-free assays comprising the membrane-bound form of the polypeptide, it may be desirable to utilize a solubilizing agent such that the membrane-bound form of the polypeptide is maintained in solution. Examples of such solubilizing agents include but are not limited to non-ionic detergents such as n-octylglucoside, n-dodecylglucoside, n-dodecylmaltoside, octanoyl-N-methylglucamide, decanoyl-N-methylglucamide, Triton X-100, Triton X-114, Thesit, Isotridecypoly(ethylene glycol ether)_n, 3-[(3-cholamidopropyl)dimethylamminio]-1-propane sulfonate (CHAPS), 3-[(3-cholamidopropyl)-dimethylamminio]-2-hydroxy-1-propane sulfonate (CHAPSO), or N-dodecyl=N,N-dimethyl-3-ammonio-1-propane sulfonate.

In various embodiments of the above assay methods of the present invention, it may be desirable to immobilize KCNK1 (or a KCNK1 target molecule) to facilitate separation of complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay. Binding of a test compound to KCNK1, or interaction of KCNK1 with a target molecule in the presence and absence of a candidate compound, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtitre plates, test tubes, and micro-centrifuge tubes. In one embodiment, a fusion protein can be provided which adds a domain that allows one or both of the proteins to be bound to a matrix. For example, glutathione-S-transferase (GST) fusion proteins or glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical; St. Louis, Mo.) or glutathione derivatized microtitre plates, which are then combined with the test compound or the test compound and either the non-adsorbed target protein or KCNK1, and the mixture incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads or microtitre plate wells are washed to remove any unbound components and complex formation is measured either directly or indirectly, for example, as described above. Alternatively, the complexes can be dissociated from the matrix, and the level of binding or activity of KCNK1 can be determined using standard techniques.

Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, either KCNK1 or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated polypeptide of the invention or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well known in the art (e.g., biotinylation kit, Pierce Chemicals; Rockford, Ill.), and immobilized in the wells of streptavidin-coated plates (Pierce Chemical). Alternatively, antibodies reactive with KCNK1 or target molecules but which do not interfere with binding of the polypeptide of the invention to its target molecule can be derivatized to the wells of the plate, and unbound target or polypeptide of the invention trapped in the wells by antibody conjugation. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with KCNK1 or target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with KCNK1 or target molecule.

The screening assay can also involve monitoring the expression of KCNK1. For example, regulators of expression of KCNK1 can be identified in a method in which a cell is contacted with a candidate compound and the expression of KCNK1 protein or mRNA in the cell is determined. The level of expression of KCNK1 protein or mRNA the presence of the candidate compound is compared to the level of expression of KCNK1 protein or mRNA in the absence of the candidate

compound. The candidate compound can then be identified as a regulator of expression of KCNK1 based on this comparison. For example, when expression of KCNK1 protein or mRNA protein is greater (statistically significantly greater) in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of KCNK1 protein or mRNA expression. Alternatively, when expression of KCNK1 protein or mRNA is less (statistically significantly less) in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of KCNK1 protein or mRNA expression. The level of KCNK1 protein or mRNA expression in the cells can be determined by methods described below.

10 *Binding Assays*

For binding assays, the test compound is preferably a small molecule which potentially binds to a regulatory channel site or nearby the channel pore of KCNK1, thereby competing with physiological modulators for said regulatory site, blocking the channel pore, or holding it in its open state such that normal biological activity is prevented, enhanced or otherwise modified. Examples of such small molecules include, but are not limited to, small peptides or peptide-like molecules as well as small compounds.. Potential ligands which bind to a polypeptide of the invention include, but are not limited to, the natural ligands of known KCNK1 ion channels and analogues or derivatives thereof.

In binding assays, either the test compound or the KCNK1 polypeptide can comprise a detectable label, such as a fluorescent, radioisotopic, chemiluminescent, or enzymatic label, such as horseradish peroxidase, alkaline phosphatase, or luciferase. Detection of a test compound which is bound to KCNK1 polypeptide can then be accomplished, for example, by direct counting of radioemission, by scintillation counting, or by determining conversion of an appropriate substrate to a detectable product. Alternatively, binding of a test compound to a KCNK1 polypeptide can be determined without labeling either of the interactants. For example, a microphysiometer can be used to detect binding of a test compound with a KCNK1 polypeptide. A microphysiometer (e.g., Cytosensor™) is an analytical instrument that measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate can be used as an indicator of the interaction between a test compound and KCNK1 [Haseloff, (1988)].

Determining the ability of a test compound to bind to KCNK1 also can be accomplished using a technology such as real-time Bimolecular Interaction Analysis (BIA) [McConnell, (1992); Sjolander, (1991)]. BIA is a technology for studying biospecific interactions in real time, without labeling any of the interactants (e.g., BIAcore™). Changes in the optical phenomenon surface

plasmon resonance (SPR) can be used as an indication of real-time reactions between biological molecules.

In a competitive binding format, the assay includes contacting KCNK1 with a known compound which carries a detectable label and which is coupled to a detectable label and which binds
5 KCNK1 to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with KCNK1, wherein determining the ability of the test compound to interact with KCNK1 comprises determining the ability of the test compound to preferentially bind to KCNK1 as compared to the known compound.

In yet another aspect of the invention, a KCNK1-like polypeptide can be used as a "bait protein" in
10 a two-hybrid assay or three-hybrid assay [Szabo, (1995); U.S. 5,283,317], to identify other proteins which bind to or interact with KCNK1 and modulate its activity.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. For example, in one construct, polynucleotide encoding KCNK1 can be fused to a
15 polynucleotide encoding the DNA binding domain of a known transcription factor (*e.g.*, GAL-4). In the other construct a DNA sequence that encodes an unidentified protein ("prey" or "sample") can be fused to a polynucleotide that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact *in vivo* to form a protein-dependent complex, the DNA-binding and activation domains of the transcription factor are
20 brought into close proximity. This proximity allows transcription of a reporter gene (*e.g.*, LacZ), which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected, and cell colonies containing the functional transcription factor can be isolated and used to obtain the DNA sequence encoding the protein which interacts with KCNK1.

25 It may be desirable to immobilize either the KCNK1 (or polynucleotide) or the test compound to facilitate separation of the bound form from unbound forms of one or both of the interactants, as well as to accommodate automation of the assay. Thus, either the KCNK1-like polypeptide (or polynucleotide) or the test compound can be bound to a solid support. Suitable solid supports include, but are not limited to, glass or plastic slides, tissue culture plates, microtiter wells, tubes,
30 silicon chips, or particles such as beads (including, but not limited to, latex, polystyrene, or glass beads). Any method known in the art can be used to attach KCNK1-like polypeptide (or polynucleotide) or test compound to a solid support, including use of covalent and non-covalent linkages, passive absorption, or pairs of binding moieties attached respectively to the polypeptide (or polynucleotide) or test compound and the solid support. Test compounds are preferably bound

to the solid support in an array, so that the location of individual test compounds can be tracked. Binding of a test compound to KCNK1 (or a polynucleotide encoding for KCNK1) can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and microcentrifuge tubes.

- 5 In one embodiment, KCNK1 is a fusion protein comprising a domain that allows binding of KCNK1 to a solid support. For example, glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, Mo.) or glutathione derivatized microtiter plates, which are then combined with the test compound or the test compound and the non-adsorbed KCNK1; the mixture is then incubated under conditions
- 10 conducive to complex formation (*e.g.*, at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components. Binding of the interactants can be determined either directly or indirectly, as described above. Alternatively, the complexes can be dissociated from the solid support before binding is determined.
- 15 Other techniques for immobilizing proteins or polynucleotides on a solid support also can be used in the screening assays of the invention. For example, either KCNK1 (or a polynucleotide encoding KCNK1) or a test compound can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated KCNK1 (or a polynucleotide encoding biotinylated KCNK1) or test compounds can be prepared from biotin-NHS (N-hydroxysuccinimide) using techniques well
- 20 known in the art (*e.g.*, biotinylation kit, Pierce Chemicals, Rockford, Ill.) and immobilized in the wells of streptavidin-coated plates (Pierce Chemical). Alternatively, antibodies which specifically bind to KCNK1, polynucleotide, or a test compound, but which do not interfere with a desired binding site, such as the active site of KCNK1, can be derivatized to the wells of the plate. Unbound target or protein can be trapped in the wells by antibody conjugation.
- 25 Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies which specifically bind to KCNK1 polypeptide or test compound, enzyme-linked assays which rely on detecting an activity of KCNK1 polypeptide, and SDS gel electrophoresis under non-reducing conditions.
- 30 Screening for test compounds which bind to a KCNK1 polypeptide or polynucleotide also can be carried out in an intact cell. Any cell which comprises a KCNK1 polypeptide or polynucleotide can be used in a cell-based assay system. A KCNK1 polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Binding of the test compound to KCNK1 or a polynucleotide encoding KCNK1 is determined as described above.

Functional Assays

- Test compounds can be tested for the ability to increase or decrease KCNK1 functional activity of a KCNK1 polypeptide. The KCNK1 activity can be measured, for example, using methods described in the specific examples, below. KCNK1 functional activity can be measured after
- 5 contacting an intact cell having functional ion channel activity with a test compound. A test compound which decreases KCNK1 activity by at least about 10, preferably about 50, more preferably about 75, 90, or 100% is identified as a potential agent for decreasing KCNK1 activity. A test compound which increases KCNK1 activity by at least about 10, preferably about 50, more preferably about 75, 90, or 100% is identified as a potential agent for increasing KCNK1 activity.
- 10 Other screening techniques include the use of cells which express KCNK1 (for example, transfected CHO cells) in a system which measures extracellular pH changes caused by channel activation [Iwabuchi, (1993)]. For example, compounds may be contacted with a cell which expresses the channel polypeptide of the present invention and a second messenger response, *e.g.*, signal transduction or pH changes, can be measured to determine whether the potential compound
- 15 activates or inhibits the channel. Another such screening technique involves introducing RNA or DNA encoding KCNK1 into *Xenopus* oocytes to transiently express the channel. The receptor oocytes can then be contacted with the channel ligand or activated otherwise and a compound to be screened, followed by detection of modulation of the observed control signal (Schnitzler et al., 2003 Receptor Channels 9 (1), 41-48) Other screening techniques include the use of cells which
- 20 express KCNK1 (for example, transfected CHO cells) in a system which measures voltage changes caused by channel activation (Baxter et al., 2002 J Biomol Screen 7(1), 79-85) For example, compounds may be contacted with a cell which expresses the channel polypeptide of the present invention and voltage changes can be measured to determine whether the potential compound activates or inhibits the channel.
- 25 Another screening technique involves expressing KCNK1 in cells in which the channel is linked to a phospholipase C or D. Such cells include endothelial cells, smooth muscle cells, embryonic kidney cells, etc. The screening may be accomplished as described above by quantifying the degree of activation of the channel from changes in the phospholipase activity.

Gene Expression

- 30 In another embodiment, test compounds which increase or decrease KCNK1 gene expression are identified. As used herein, the term "correlates with expression of a polynucleotide" indicates that the detection of the presence of nucleic acids, the same or related to a nucleic acid sequence encoding KCNK1, by northern analysis or real-time PCR is indicative of the presence of nucleic

acids encoding KCNK1 in a sample, and thereby correlates with expression of the transcript from the polynucleotide encoding KCNK1. The term "microarray", as used herein, refers to an array of distinct polynucleotides or oligonucleotides arrayed on a substrate, such as paper, nylon or any other type of membrane, filter, chip, glass slide, or any other suitable solid support. A KCNK1 polynucleotide is contacted with a test compound, and the expression of an RNA or polypeptide product of KCNK1 polynucleotide is determined. The level of expression of appropriate mRNA or polypeptide in the presence of the test compound is compared to the level of expression of mRNA or polypeptide in the absence of the test compound. The test compound can then be identified as a regulator of expression based on this comparison. For example, when expression of mRNA or polypeptide is greater in the presence of the test compound than in its absence, the test compound is identified as a stimulator or enhancer of the mRNA or polypeptide expression. Alternatively, when expression of the mRNA or polypeptide is less in the presence of the test compound than in its absence, the test compound is identified as an inhibitor of the mRNA or polypeptide expression.

The level of KCNK1 mRNA or polypeptide expression in the cells can be determined by methods well known in the art for detecting mRNA or polypeptide. Either qualitative or quantitative methods can be used. The presence of polypeptide products of KCNK1 polynucleotide can be determined, for example, using a variety of techniques known in the art, including immunochemical methods such as radioimmunoassay, Western blotting, and immunohistochemistry. Alternatively, polypeptide synthesis can be determined *in vivo*, in a cell culture, or in an *in vitro* translation system by detecting incorporation of labelled amino acids into KCNK1.

Such screening can be carried out either in a cell-free assay system or in an intact cell. Any cell which expresses KCNK1 polynucleotide can be used in a cell-based assay system. The KCNK1 polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Either a primary culture or an established cell line can be used.

Test Compounds

Suitable test compounds for use in the screening assays of the invention can be obtained from any suitable source, e.g., conventional compound libraries. The test compounds can also be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the "one-bead one-compound" library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds [Lam, (1997)]. Examples of

methods for the synthesis of molecular libraries can be found in the art. Libraries of compounds may be presented in solution or on beads, bacteria, spores, plasmids or phage.

Modeling of Regulators

Computer modeling and searching technologies permit identification of compounds, or the
5 improvement of already identified compounds, that can modulate KCNK1 expression or activity. Having identified such a compound or composition, the active sites or regions are identified. Such active sites might typically be ligand binding sites, such as the interaction domain of the ligand with KCNK1. The active site can be identified using methods known in the art including, for example, from the amino acid sequences of peptides, from the nucleotide sequences of nucleic
10 acids, or from study of complexes of the relevant compound or composition with its natural ligand. In the latter case, chemical or X-ray crystallographic methods can be used to find the active site by finding where on the factor the complexed ligand is found.

Next, the three dimensional geometric structure of the active site is determined. This can be done by known methods, including X-ray crystallography, which can determine a complete molecular
15 structure. On the other hand, solid or liquid phase NMR can be used to determine certain intramolecular distances. Any other experimental method of structure determination can be used to obtain partial or complete geometric structures. The geometric structures may be measured with a complexed ligand, natural or artificial, which may increase the accuracy of the active site structure determined.

20 If an incomplete or insufficiently accurate structure is determined, the methods of computer based numerical modeling can be used to complete the structure or improve its accuracy. Any recognized modeling method may be used, including parameterized models specific to particular biopolymers such as proteins or nucleic acids, molecular dynamics models based on computing molecular motions, statistical mechanics models based on thermal ensembles, or combined models.
25 For most types of models, standard molecular force fields, representing the forces between constituent atoms and groups, are necessary, and can be selected from force fields known in physical chemistry. The incomplete or less accurate experimental structures can serve as constraints on the complete and more accurate structures computed by these modeling methods.

Finally, having determined the structure of the active site, either experimentally, by modeling, or
30 by a combination, candidate modulating compounds can be identified by searching databases containing compounds along with information on their molecular structure. Such a search seeks compounds having structures that match the determined active site structure and that interact with

the groups defining the active site. Such a search can be manual, but is preferably computer assisted. These compounds found from this search are potential KCNK1 modulating compounds.

Alternatively, these methods can be used to identify improved modulating compounds from an already known modulating compound or ligand. The composition of the known compound can be
5 modified and the structural effects of modification can be determined using the experimental and computer modeling methods described above applied to the new composition. The altered structure is then compared to the active site structure of the compound to determine if an improved fit or interaction results. In this manner systematic variations in composition, such as by varying side groups, can be quickly evaluated to obtain modified modulating compounds or ligands of
10 improved specificity or activity.

Therapeutic Indications and Methods

It was found by the present applicant that KCNK1 is expressed in various human tissues.

Neurology

CNS disorders include disorders of the central nervous system as well as disorders of the
15 peripheral nervous system.

CNS disorders include, but are not limited to brain injuries, cerebrovascular diseases and their consequences, Parkinson's disease, corticobasal degeneration, motor neuron disease, dementia, including ALS, multiple sclerosis, traumatic brain injury, stroke, post-stroke, post-traumatic brain injury, and small-vessel cerebrovascular disease. Dementias, such as Alzheimer's disease,
20 vascular dementia, dementia with Lewy bodies, frontotemporal dementia and Parkinsonism linked to chromosome 17, frontotemporal dementias, including Pick's disease, progressive nuclear palsy, corticobasal degeneration, Huntington's disease, thalamic degeneration, Creutzfeld-Jakob dementia, HIV dementia, schizophrenia with dementia, and Korsakoff's psychosis, within the meaning of the definition are also considered to be CNS disorders.

25 Similarly, cognitive-related disorders, such as mild cognitive impairment, age-associated memory impairment, age-related cognitive decline, vascular cognitive impairment, attention deficit disorders, attention deficit hyperactivity disorders, and memory disturbances in children with learning disabilities are also considered to be CNS disorders.

Pain, within the meaning of this definition, is also considered to be a CNS disorder. Pain can be
30 associated with CNS disorders, such as multiple sclerosis, spinal cord injury, sciatica, failed back surgery syndrome, traumatic brain injury, epilepsy, Parkinson's disease, post-stroke, and vascular

lesions in the brain and spinal cord (e.g., infarct, hemorrhage, vascular malformation). Non-central neuropathic pain includes that associated with post mastectomy pain, phantom feeling, reflex sympathetic dystrophy (RSD), trigeminal neuralgia, radioculopathy, post-surgical pain, HIV/AIDS related pain, cancer pain, metabolic neuropathies (e.g., diabetic neuropathy, vasculitic neuropathy secondary to connective tissue disease), paraneoplastic polyneuropathy associated, for example, with carcinoma of lung, or leukemia, or lymphoma, or carcinoma of prostate, colon or stomach, trigeminal neuralgia, cranial neuralgias, and post-herpetic neuralgia. Pain associated with peripheral nerve damage, central pain (i.e. due to cerebral ischemia) and various chronic pain i.e., lumbago, back pain (low back pain), inflammatory and/or rheumatic pain. Headache pain (for example, migraine with aura, migraine without aura, and other migraine disorders), episodic and chronic tension-type headache, tension-type like headache, cluster headache, and chronic paroxysmal hemicrania are also CNS disorders.

Visceral pain such as pancreatitis, intestinal cystitis, dysmenorrhea, irritable Bowel syndrome, Crohn's disease, biliary colic, ureteral colic, myocardial infarction and pain syndromes of the pelvic cavity, e.g., vulvodynia, orchialgia, urethral syndrome and protatodynia are also CNS disorders.

Also considered to be a disorder of the nervous system are acute pain, for example postoperative pain, and pain after trauma.

The human KCNK1 is highly expressed in the following brain tissues: brain, Alzheimer brain, cerebellum, cerebellum (right), cerebellum (left), cerebral cortex, Alzheimer cerebral cortex, frontal lobe, occipital lobe, parietal lobe, temporal lobe, precentral gyrus, postcentral gyrus, tonsilla cerebelli, vermis cerebelli, pons, cerebral peduncles, corpus callosum, hippocampus, thalamus, spinal cord, neuroblastoma IMR32 cells. The expression in brain tissues and in particular the differential expression between diseased tissue Alzheimer brain and healthy tissue brain, between diseased tissue Alzheimer cerebral cortex and healthy tissue cerebral cortex demonstrates that the human KCNK1 or mRNA can be utilized to diagnose nervous system diseases. Additionally the activity of the human KCNK1 can be modulated to treat nervous system diseases.

Cardiovascular Disorders

Heart failure is defined as a pathophysiological state in which an abnormality of cardiac function is responsible for the failure of the heart to pump blood at a rate commensurate with the requirement of the metabolizing tissue. It includes all forms of pumping failures such as high-output and low-

output, acute and chronic, right-sided or left-sided, systolic or diastolic, independent of the underlying cause.

Myocardial infarction (MI) is generally caused by an abrupt decrease in coronary blood flow that follows a thrombotic occlusion of a coronary artery previously narrowed by arteriosclerosis. MI
5 prophylaxis (primary and secondary prevention) is included as well as the acute treatment of MI and the prevention of complications.

Ischemic diseases are conditions in which the coronary flow is restricted resulting in a perfusion which is inadequate to meet the myocardial requirement for oxygen. This group of diseases includes stable angina, unstable angina and asymptomatic ischemia.

10 Arrhythmias include all forms of atrial and ventricular tachyarrhythmias, atrial tachycardia, atrial flutter, atrial fibrillation, atrio-ventricular reentrant tachycardia, preexcitation syndrome, ventricular tachycardia, ventricular flutter, ventricular fibrillation, as well as bradycardic forms of arrhythmias.

Hypertensive vascular diseases include primary as well as all kinds of secondary arterial hyper-
15 tension, renal, endocrine, neurogenic, others. The genes may be used as drug targets for the treatment of hypertension as well as for the prevention of all complications arising from cardiovascular diseases.

Peripheral vascular diseases are defined as vascular diseases in which arterial and/or venous flow is reduced resulting in an imbalance between blood supply and tissue oxygen demand. It includes
20 chronic peripheral arterial occlusive disease (PAOD), acute arterial thrombosis and embolism, inflammatory vascular disorders, Raynaud's phenomenon and venous disorders.

Atherosclerosis is a cardiovascular disease in which the vessel wall is remodeled, compromising the lumen of the vessel. The atherosclerotic remodeling process involves accumulation of cells, both smooth muscle cells and monocyte/macrophage inflammatory cells, in the intima of the vessel
25 wall. These cells take up lipid, likely from the circulation, to form a mature atherosclerotic lesion. Although the formation of these lesions is a chronic process, occurring over decades of an adult human life, the majority of the morbidity associated with atherosclerosis occurs when a lesion ruptures, releasing thrombogenic debris that rapidly occludes the artery. When such an acute event occurs in the coronary artery, myocardial infarction can ensue, and in the worst case, can result in
30 death.

The formation of the atherosclerotic lesion can be considered to occur in five overlapping stages such as migration, lipid accumulation, recruitment of inflammatory cells, proliferation of vascular

smooth muscle cells, and extracellular matrix deposition. Each of these processes can be shown to occur in man and in animal models of atherosclerosis, but the relative contribution of each to the pathology and clinical significance of the lesion is unclear.

Thus, a need exists for therapeutic methods and agents to treat cardiovascular pathologies, such as
5 atherosclerosis and other conditions related to coronary artery disease.

Cardiovascular diseases include but are not limited to disorders of the heart and the vascular system like congestive heart failure, myocardial infarction, ischemic diseases of the heart, all kinds of atrial and ventricular arrhythmias, hypertensive vascular diseases, peripheral vascular diseases, and atherosclerosis.

10 Too high or too low levels of fats in the bloodstream, especially cholesterol, can cause long-term problems. The risk to develop atherosclerosis and coronary artery or carotid artery disease (and thus the risk of having a heart attack or stroke) increases with the total cholesterol level increasing. Nevertheless, extremely low cholesterol levels may not be healthy. Examples of disorders of lipid metabolism are hyperlipidemia (abnormally high levels of fats (cholesterol, triglycerides, or both)
15 in the blood, may be caused by family history of hyperlipidemia), obesity, a high-fat diet, lack of exercise, moderate to high alcohol consumption, cigarette smoking, poorly controlled diabetes, and an underactive thyroid gland), hereditary hyperlipidemias (type I hyperlipoproteinemia (familial hyperchylomicronemia), type II hyperlipoproteinemia (familial hypercholesterolemia), type III hyperlipoproteinemia, type IV hyperlipoproteinemia, or type V hyperlipoproteinemia),
20 hypolipoproteinemia, lipidoses (caused by abnormalities in the enzymes that metabolize fats), Gaucher's disease, Niemann-Pick disease, Fabry's disease, Wolman's disease, cerebrotendinous xanthomatosis, sitosterolemia, Refsum's disease, or Tay-Sachs disease.

Kidney disorders may lead to hypertension or hypotension. Examples for kidney problems possibly leading to hypertension are renal artery stenosis, pyelonephritis, glomerulonephritis, kidney
25 tumors, polycystic kidney disease, injury to the kidney, or radiation therapy affecting the kidney. Excessive urination may lead to hypotension.

The human KCNK1 is highly expressed in the following cardiovascular related tissues: heart, heart atrium (right), heart atrium (left), interventricular septum, aorta, thrombocytes. Expression in the above mentioned tissues demonstrates that the human KCNK1 or mRNA can be utilized to
30 diagnose of cardiovascular diseases. Additionally the activity of the human KCNK1 can be modulated to treat cardiovascular diseases.

Hematological Disorders

Hematological disorders comprise diseases of the blood and all its constituents as well as diseases of organs and tissues involved in the generation or degradation of all the constituents of the blood. They include but are not limited to 1) Anemias, 2) Myeloproliferative Disorders, 3) Hemorrhagic Disorders, 4) Leukopenia, 5) Eosinophilic Disorders, 6) Leukemias, 7) Lymphomas, 8) Plasma Cell Dyscrasias, 9) Disorders of the Spleen in the course of hematological disorders. Disorders according to 1) include, but are not limited to anemias due to defective or deficient hem synthesis, deficient erythropoiesis. Disorders according to 2) include, but are not limited to polycythemia vera, tumor-associated erythrocytosis, myelofibrosis, thrombocythemia. Disorders according to 3) include, but are not limited to vasculitis, thrombocytopenia, heparin-induced thrombocytopenia, thrombotic thrombocytopenic purpura, hemolytic-uremic syndrome, hereditary and acquired disorders of platelet function, hereditary coagulation disorders. Disorders according to 4) include, but are not limited to neutropenia, lymphocytopenia. Disorders according to 5) include, but are not limited to hypereosinophilia, idiopathic hypereosinophilic syndrome. Disorders according to 6) include, but are not limited to acute myeloic leukemia, acute lymphoblastic leukemia, chronic myelocytic leukemia, chronic lymphocytic leukemia, myelodysplastic syndrome. Disorders according to 7) include, but are not limited to Hodgkin's disease, non-Hodgkin's lymphoma, Burkitt's lymphoma, mycosis fungoides cutaneous T-cell lymphoma. Disorders according to 8) include, but are not limited to multiple myeloma, macroglobulinemia, heavy chain diseases. In extension of the preceding idiopathic thrombocytopenic purpura, iron deficiency anemia, megaloblastic anemia (vitamin B12 deficiency), aplastic anemia, thalassemia, malignant lymphoma bone marrow invasion, malignant lymphoma skin invasion, hemolytic uremic syndrome, giant platelet disease are considered to be hematological diseases too.

The human KCNK1 is highly expressed in the following tissues of the hematological system: Jurkat (T-cells), thrombocytes, bone marrow CD71+ cells, bone marrow CD15+ cells, cord blood CD71+ cells. The expression in the above mentioned tissues demonstrates that the human KCNK1 or mRNA can be utilized to diagnose of hematological diseases. Additionally the activity of the human KCNK1 can be modulated to treat hematological disorders.

Gastrointestinal and Liver Diseases

Gastrointestinal diseases comprise primary or secondary, acute or chronic diseases of the organs of the gastrointestinal tract which may be acquired or inherited, benign or malignant or metaplastic, and which may affect the organs of the gastrointestinal tract or the body as a whole. They comprise but are not limited to 1) disorders of the esophagus like achalasia, vigorous achalasia, dysphagia, cricopharyngeal incoordination, pre-esophageal dysphagia, diffuse esophageal spasm, globus

sensation, Barrett's metaplasia, gastroesophageal reflux, 2) disorders of the stomach and duodenum like functional dyspepsia, inflammation of the gastric mucosa, gastritis, stress gastritis, chronic erosive gastritis, atrophy of gastric glands, metaplasia of gastric tissues, gastric ulcers, duodenal ulcers, neoplasms of the stomach, 3) disorders of the pancreas like acute or chronic pancreatitis, insufficiency of the exocrine or endocrine tissues of the pancreas like steatorrhea, diabetes, neoplasms of the exocrine or endocrine pancreas like 3.1) multiple endocrine neoplasia syndrome, ductal adenocarcinoma, cystadenocarcinoma, islet cell tumors, insulinoma, gastrinoma, carcinoid tumors, glucagonoma, Zollinger-Ellison syndrome, Vipoma syndrome, malabsorption syndrome, 4) disorders of the bowel like chronic inflammatory diseases of the bowel, Crohn's disease, ileus, diarrhea and constipation, colonic inertia, megacolon, malabsorption syndrome, ulcerative colitis, 4.1) functional bowel disorders like irritable bowel syndrome, 4.2) neoplasms of the bowel like familial polyposis, adenocarcinoma, primary malignant lymphoma, carcinoid tumors, Kaposi's sarcoma, polyps, cancer of the colon and rectum.

Liver diseases comprise primary or secondary, acute or chronic diseases or injury of the liver which may be acquired or inherited, benign or malignant, and which may affect the liver or the body as a whole. They comprise but are not limited to disorders of the bilirubin metabolism, jaundice, syndromes of Gilbert's, Crigler-Najjar, Dubin-Johnson and Rotor; intrahepatic cholestasis, hepatomegaly, portal hypertension, ascites, Budd-Chiari syndrome, portal-systemic encephalopathy, fatty liver, steatosis, Reye's syndrome, liver diseases due to alcohol, alcoholic hepatitis or cirrhosis, fibrosis and cirrhosis, fibrosis and cirrhosis of the liver due to inborn errors of metabolism or exogenous substances, storage diseases, syndromes of Gaucher's, Zellweger's, Wilson's - disease, acute or chronic hepatitis, viral hepatitis and its variants, inflammatory conditions of the liver due to viruses, bacteria, fungi, protozoa, helminths; drug induced disorders of the liver, chronic liver diseases like primary sclerosing cholangitis, α_1 -antitrypsin-deficiency, primary biliary cirrhosis, postoperative liver disorders like postoperative intrahepatic cholestasis, hepatic granulomas, vascular liver disorders associated with systemic disease, benign or malignant neoplasms of the liver, disturbance of liver metabolism in the new-born or prematurely born.

The human KCNK1 is highly expressed in the following tissues of the gastroenterological system: esophagus, esophagus tumor, colon, colon tumor, ileum, ileum tumor, rectum, salivary gland. The expression in the above mentioned tissues demonstrates that the human KCNK1 or mRNA can be utilized to diagnose of gastroenterological disorders. Additionally the activity of the human KCNK1 can be modulated to treat gastroenterological disorders.

Dermatologic Disorders

The skin serves several functions. It's an multi-layered organ system that builds an effective protective cover and regulates body temperature, senses painful and pleasant stimuli, keeps substances from entering the body, and provides a shield from the sun's harmful effects. Skin color, texture, and folds help mark people as individuals. Thus, skin disorders or diseases often have important consequences for physical and mental health. Skin disorders include, but are not limited to the conditions described in the following.

Itching (pruritus) is a sensation that instinctively demands scratching, which may be caused by a skin condition or a systemic diseases.

Superficial Skin Disorders affect the uppermost layer of the skin, the stratum corneum or the keratin layer, and it consists of many layers of flattened, dead cells and acts as a barrier to protect the underlying tissue from injury and infection. Disorders of the superficial skin layers involve the stratum corneum and deeper layers of the epidermis.

Examples of superficial skin disorders are provided in the following.

Dry skin often occurs in people past middle age, severe dry skin (ichthyosis) results from an inherited scaling disease, such as ichthyosis vulgaris or epidermolytic hyperkeratosis. Ichthyosis also results from nonhereditary disorders, such as leprosy, underactive thyroid, lymphoma, AIDS, and sarcoidosis.

Keratosis pilaris is a common disorder in which dead cells shed from the upper layer of skin and form plugs that fill the openings of hair follicles.

A callus is an area on the stratum corneum or keratin layer, that becomes abnormally thick in response to repeated rubbing.

A corn is a pea-sized, thickened area of keratin that occurs on the feet.

Psoriasis is a chronic, recurring disease recognizable by silvery scaling bumps and various-sized plaques (raised patches). An abnormally high rate of growth and turnover of skin cells causes the scaling.

Pityriasis rosea is a mild disease that causes scaly, rose-colored, inflamed skin. Pityriasis rosea is possibly caused by an infectious agent, although none has been identified.

Lichen planus, a recurring itchy disease, starts as a rash of small discrete bumps that then combine and become rough, scaly plaques (raised patches).

Dermatitis (eczema) is an inflammation of the upper layers of the skin, causing blisters, redness, swelling, oozing, scabbing, scaling, and usually itching.

- 5 Forms of dermatitis are contact dermatitis, or chronic dermatitis of the hands and feet, e.g. Pompholyx.

Further examples of dermatitic disorders are atopic dermatitis, seborrheic dermatitis, nummular dermatitis, generalized exfoliative dermatitis, stasis dermatitis, or localized scratch dermatitis (lichen simplex chronicus, neurodermatitis).

- 10 Other skin disorders are caused by inflammation. The skin can break out in a variety of rashes, sores, and blisters. Some skin eruptions can even be life threatening.

Drug rashes are side effects of medications, mainly allergic reactions to medications.

- 15 Toxic epidermal necrolysis is a life-threatening skin disease in which the top layer of the skin peels off in sheets. This condition can be caused by a reaction to a drug, or by some other serious disease.

Erythema multiforme, often caused by herpes simplex is a disorder characterized by patches of red, raised skin that often look like targets and usually are distributed symmetrically over the body.

Erythema nodosum is an inflammatory disorder that produces tender red bumps (nodules) under the skin, most often over the shins but occasionally on the arms and other areas.

- 20 Granuloma annulare is a chronic skin condition of unknown cause in which small, firm, raised bumps form a ring with normal or slightly sunken skin in the center.

Some skin disorders are characterized as blistering diseases. Three autoimmune diseases-- pemphigus, bullous pemphigoid, and dermatitis herpetiformis--are among the most serious.

- 25 Pemphigus is an uncommon, sometimes fatal, disease in which blisters (bullae) of varying sizes break out on the skin, the lining of the mouth, and other mucous membranes.

Bullous pemphigoid is an autoimmune disease that causes blistering.

Dermatitis herpetiformis is an autoimmune disease in which clusters of intensely itchy, small blisters and hive-like swellings break out and persist. In people with the disease, proteins in wheat,

rye, barley, and oat products activate the immune system, which attacks parts of the skin and somehow causes the rash and itching.

Sweating disorders also belong to skin disorders.

Prickly heat is an itchy skin rash caused by trapped sweat.

- 5 Excessive sweating (hyperhidrosis) may affect the entire surface of the skin, but often it's limited to the palms, soles, armpits, or groin. The affected area is often pink or bluish white, and in severe cases the skin may be cracked, scaly, and soft, especially on the feet.

Skin disorders can affect the sebaceous glands. The sebaceous glands, which secrete oil onto the skin, lie in the dermis, the skin layer just below the surface layer (epidermis). Sebaceous gland
10 disorders include acne, rosacea, perioral dermatitis, and sebaceous cysts.

Acne is a common skin condition in which the skin pores become clogged, leading to pimples and inflamed, infected abscesses (collections of pus). Acne tends to develop in teenagers.

Acne is further subdivided in superficial acne or deep acne.

Rosacea is a persistent skin disorder that produces redness, tiny pimples, and broken blood vessels,
15 usually on the central area of the face.

Perioral dermatitis is a red, often bumpy rash around the mouth and on the chin.

A sebaceous cyst (keratinous cyst) is a slow-growing bump containing dead skin, skin excretions, and other skin particles. These cysts may be small and can appear anywhere.

Hair Disorders also are skin disorders. Hair disorders include excessive hairiness, baldness, and
20 ingrown beard hairs.

The skin can be infected by bacteria. Bacterial skin infections can range in seriousness from minor acne to a life-threatening condition, such as staphylococcal scalded skin syndrome. The most common bacterial skin infections are caused by Staphylococcus and Streptococcus. Risk factors for skin infections are for example diabetes, AIDS or skin lesions.

- 25 Impetigo is a skin infection, caused by Staphylococcus or Streptococcus, leading to the formation of small pus-filled blisters (pustules).

Folliculitis is an inflammation of the hair follicles caused by infection with Staphylococcus. The infection damages the hairs, which can be easily pulled out.

Boils (furuncles) are large, tender, swollen, raised areas caused by staphylococcal infection around hair follicles.

Carbuncles are clusters of boils that result in extensive sloughing of skin and scar formation. Carbuncles develop and heal more slowly than single boils and may lead to fever and fatigue.

- 5 Erysipelas is a skin infection caused by *Streptococcus*. A shiny, red, slightly swollen, tender rash develops, often with small blisters. Lymph nodes around the infected area may become enlarged and painful.

- Cellulitis is a spreading infection in, and sometimes beneath, the deep layers of the skin. Cellulitis most often results from a streptococcal infection or a staphylococcal infection. However, many
10 other bacteria can also cause cellulitis.

Paronychia is an infection around the edge of a fingernail or toenail. Paronychia can be caused by many different bacteria, including *Pseudomonas* and *Proteus*, and by fungi, such as *Candida*.

- Staphylococcal scalded skin syndrome is a widespread skin infection that can lead to toxic shock syndrome, in which the skin peels off as though burned. Certain types of staphylococci produce a
15 toxic substance that causes the top layer of skin (epidermis) to split from the rest of the skin.

Erythrasma is an infection of the top layers of the skin by the bacterium *Corynebacterium minutissimum*.

- Skin infections are often caused by fungi. Fungi that infect the skin (dermatophytes) live only in the dead, topmost layer (stratum corneum) and don't penetrate deeper. Some fungal infections
20 cause no symptoms or produce only a small amount of irritation, scaling, and redness. Other fungal infections cause itching, swelling, blisters, and severe scaling.

Ringworm is a fungal skin infection caused by several different fungi and generally classified by its location on the body.

- Examples are Athlete's foot (foot ringworm, caused by either *Trichophyton* or *Epidermophyton*),
25 jock itch (groin ringworm, can be caused by a variety of fungi and yeasts), scalp ringworm, caused by *Trichophyton* or *Microsporum*), nail ringworm and body ringworm (caused by *Trichophyton*).

- Candidiasis (yeast infection, moniliasis) is an infection by the yeast *Candida*. *Candida* usually infects the skin and mucous membranes, such as the lining of the mouth and vagina. Rarely, it invades deeper tissues as well as the blood, causing life-threatening systemic candidiasis. The
30 following types of candida infections can be distinguished: Infections in skinfolds (intertriginous

infections), vaginal and penile candida infections (vulvovaginitis), thrush, Perlèche (candida infection at the corners of the mouth), candidal paronychia (candida growing in the nail beds, produces painful swelling and pus).

Tinea versicolor is a fungal infection that causes white to light brown patches on the skin.

- 5 The skin can also be affected by parasites, mainly tiny insects or worms.

Scabies is a mite infestation that produces tiny reddish pimples and severe itching. Scabies is caused by the itch mite *Sarcoptes scabiei*.

Lice infestation (pediculosis) causes intense itching and can affect almost any area of the skin. Head lice and pubic lice are two different species.

- 10 Creeping eruption (cutaneous larva migrans) is a hookworm infection transmitted from warm, moist soil to exposed skin. The infection is caused by a hookworm that normally inhabits dogs and cats.

- Many types of viruses invade the skin. The medically important ones cause warts and cold sores (fever blisters) on the lip. Warts are caused by the papillomavirus, and cold sores are caused by the herpes simplex virus. Another important group of viruses that infect the skin belongs to the poxvirus family. Chickenpox remains a common childhood infection. A poxvirus also causes molluscum contagiosum, which is an infection of the skin by a poxvirus that causes skin-colored, smooth, waxy bumps.
- 15

- Sunlight can cause severe skin damage. Sunburn results from an overexposure to ultraviolet B (UVB) rays. Some sunburned people develop a fever, chills, and weakness, and those with very bad sunburns even may go into shock--low blood pressure, and fainting.
- 20

People who are in the sun a lot have an increased risk of skin cancers, including squamous cell carcinoma, basal cell carcinoma, and to some degree, malignant melanoma.

- Drugs, among other causes, can cause skin photosensitivity reactions which can occur after only a few minutes of sun exposure. These reactions include redness, peeling, hives, blisters, and thickened, scaling patches (photosensitivity).
- 25

Some skin disorders are characterized as Pigment Disorders.

Albinism is a rare, inherited disorder in which no melanin is formed.

Vitiligo is a condition in which a loss of melanocytes results in smooth, whitish patches of skin, which may occur after unusual physical trauma and tends to occur with certain other diseases, including Addison's disease, diabetes, pernicious anemia, and thyroid disease.

Tinea versicolor is a fungal infection of the skin that sometimes results in hyperpigmentation.

- 5 Melasma appears on the face (usually the forehead, cheeks, temples, and jaws) as a roughly symmetric group of dark brown patches of pigmentation that are often clearly delineated.

Skin growths, which are abnormal accumulations of different types of cells, may be present at birth or develop later. Noncancerous (benign) growth and cancerous (malignant) growth types are distinguished.

- 10 Moles (nevi) are small, usually dark, skin growths that develop from pigment-producing cells in the skin (melanocytes). Most moles are harmless. However, noncancerous moles can develop into malignant melanoma.

Skin tags are soft, small, flesh-colored or slightly darker skin flaps that appear mostly on the neck, in the armpits, or in the groin.

- 15 Lipomas are soft deposits of fatty material that grow under the skin, causing round or oval lumps.

Angiomas are collections of abnormally dense blood or lymph vessels that are usually located in and below the skin and that cause red or purple discolorations.

Examples of angiomas are port-wine stains, strawberry marks, cavernous hemangiomas, spider angiomas, and lymphangiomas.

- 20 Pyogenic granulomas are scarlet, brown, or blue-black slightly raised areas caused by increased growth of capillaries (the smallest blood vessels) and swelling of the surrounding tissue.

Seborrheic keratoses (sometimes called seborrheic warts) are flesh-colored, brown, or black growths that can appear anywhere on the skin.

- 25 Dermatofibromas are small, red-to-brown bumps (nodules) that result from an accumulation of fibroblasts, the cells that populate the soft tissue under the skin.

Keratoacanthomas are round, firm, usually flesh-colored growths that have an unusual central crater containing a pasty material.

Keloids are smooth, shiny, slightly pink, often dome-shaped, proliferative growths of fibrous tissue that form over areas of injury or over surgical wounds.

Skin cancer is the most common form of cancer, but most types of skin cancers are curable.

Basal cell carcinoma is a cancer that originates in the lowest layer of the epidermis.

5 Squamous cell carcinoma is cancer that originates in the middle layer of the epidermis.

Bowen's disease is a form of squamous cell carcinoma that's confined to the epidermis and hasn't yet invaded the underlying dermis.

Melanoma is a cancer that originates in the pigment-producing cells of the skin (melanocytes).

Kaposi's sarcoma is a cancer that originates in the blood vessels, usually of the skin.

10 Paget's disease is a rare type of skin cancer that looks like an inflamed, reddened patch of skin (dermatitis); it originates in glands in or under the skin.

The human KCNK1 is highly expressed in the following dermatological tissues: skin. The expression in the above mentioned tissues demonstrates that the human KCNK1 or mRNA can be utilized to diagnose of dermatological diseases. Additionally the activity of the human KCNK1 can
15 be modulated to treat those diseases.

Cancer Disorders

Cancer disorders within the scope of this definition comprise any disease of an organ or tissue in mammals characterized by poorly controlled or uncontrolled multiplication of normal or abnormal cells in that tissue and its effect on the body as a whole. Cancer diseases within the scope of the
20 definition comprise benign neoplasms, dysplasias, hyperplasias as well as neoplasms showing metastatic growth or any other transformations like e.g. leukoplakias which often precede a breakout of cancer. Cells and tissues are cancerous when they grow more rapidly than normal cells, displacing or spreading into the surrounding healthy tissue or any other tissues of the body described as metastatic growth, assume abnormal shapes and sizes, show changes in their
25 nucleocytoplasmatic ratio, nuclear polychromasia, and finally may cease. Cancerous cells and tissues may affect the body as a whole when causing paraneoplastic syndromes or if cancer occurs within a vital organ or tissue, normal function will be impaired or halted, with possible fatal results. The ultimate involvement of a vital organ by cancer, either primary or metastatic, may lead to the death of the mammal affected. Cancer tends to spread, and the extent of its spread is usually
30 related to an individual's chances of surviving the disease. Cancers are generally said to be in one

of three stages of growth: early, or localized, when a tumor is still confined to the tissue of origin, or primary site; direct extension, where cancer cells from the tumour have invaded adjacent tissue or have spread only to regional lymph nodes; or metastasis, in which cancer cells have migrated to distant parts of the body from the primary site, via the blood or lymph systems, and have established secondary sites of infection. Cancer is said to be malignant because of its tendency to cause death if not treated. Benign tumors usually do not cause death, although they may if they interfere with a normal body function by virtue of their location, size, or paraneoplastic side effects. Hence benign tumors fall under the definition of cancer within the scope of this definition as well. In general, cancer cells divide at a higher rate than do normal cells, but the distinction between the growth of cancerous and normal tissues is not so much the rapidity of cell division in the former as it is the partial or complete loss of growth restraint in cancer cells and their failure to differentiate into a useful, limited tissue of the type that characterizes the functional equilibrium of growth of normal tissue. Cancer tissues may express certain molecular receptors and probably are influenced by the host's susceptibility and immunity and it is known that certain cancers of the breast and prostate, for example, are considered dependent on specific hormones for their existence. The term "cancer" under the scope of the definition is not limited to simple benign neoplasia but comprises any other benign and malign neoplasia like 1) Carcinoma, 2) Sarcoma, 3) Carcinosarcoma, 4) Cancers of the blood-forming tissues, 5) tumors of nerve tissues including the brain, 6) cancer of skin cells. Cancer according to 1) occurs in epithelial tissues, which cover the outer body (the skin) and line mucous membranes and the inner cavitory structures of organs e.g. such as the breast, lung, the respiratory and gastrointestinal tracts, the endocrine glands, and the genitourinary system. Ductal or glandular elements may persist in epithelial tumors, as in adenocarcinomas like e.g. thyroid adenocarcinoma, gastric adenocarcinoma, uterine adenocarcinoma. Cancers of the pavement-cell epithelium of the skin and of certain mucous membranes, such as e.g. cancers of the tongue, lip, larynx, urinary bladder, uterine cervix, or penis, may be termed epidermoid or squamous-cell carcinomas of the respective tissues and are in the scope of the definition of cancer as well. Cancer according to 2) develops in connective tissues, including fibrous tissues, adipose (fat) tissues, muscle, blood vessels, bone, and cartilage like e.g. osteogenic sarcoma; liposarcoma, fibrosarcoma, synovial sarcoma. Cancer according to 3) is cancer that develops in both epithelial and connective tissue. Cancer disease within the scope of this definition may be primary or secondary, whereby primary indicates that the cancer originated in the tissue where it is found rather than was established as a secondary site through metastasis from another lesion. Cancers and tumor diseases within the scope of this definition may be benign or malign and may affect all anatomical structures of the body of a mammal. By example but not limited to they comprise cancers and tumor diseases of I) the bone marrow and bone marrow derived cells (leukemias), II) the endocrine and exocrine glands like e.g. thyroid, parathyroid, pituitary, adrenal

glands, salivary glands, pancreas III) the breast, like e.g. benign or malignant tumors in the mammary glands of either a male or a female, the mammary ducts, adenocarcinoma, medullary carcinoma, comedo carcinoma, Paget's disease of the nipple, inflammatory carcinoma of the young woman, IV) the lung, V) the stomach, VI) the liver and spleen, VII) the small intestine, VIII) the colon, IX) the bone and its supportive and connective tissues like malignant or benign bone tumour, e.g. malignant osteogenic sarcoma, benign osteoma, cartilage tumors; like malignant chondrosarcoma or benign chondroma; bone marrow tumors like malignant myeloma or benign eosinophilic granuloma, as well as metastatic tumors from bone tissues at other locations of the body; X) the mouth, throat, larynx, and the esophagus, XI) the urinary bladder and the internal and external organs and structures of the urogenital system of male and female like ovaries, uterus, cervix of the uterus, testes, and prostate gland, XII) the prostate, XIII) the pancreas, like ductal carcinoma of the pancreas; XIV) the lymphatic tissue like lymphomas and other tumors of lymphoid origin, XV) the skin, XVI) cancers and tumor diseases of all anatomical structures belonging to the respiration and respiratory systems including thoracal muscles and linings, XVII) primary or secondary cancer of the lymph nodes XVIII) the tongue and of the bony structures of the hard palate or sinuses, XXIV) the mouth, cheeks, neck and salivary glands, XX) the blood vessels including the heart and their linings, XXI) the smooth or skeletal muscles and their ligaments and linings, XXII) the peripheral, the autonomous, the central nervous system including the cerebellum, XXIII) the adipose tissue.

The human KCNK1 is highly expressed in the following cancer tissues: esophagus tumor, colon tumor, ileum tumor, Jurkat (T-cells), lung tumor, breast tumor. The expression in the above mentioned tissues and in particular the differential expression between diseased tissue esophagus tumor and healthy tissue esophagus, between diseased tissue colon tumor and healthy tissue colon, between diseased tissue ileum tumor and healthy tissue ileum, between diseased tissue Jurkat (T-cells) and healthy tissue, between diseased tissue lung tumor and healthy tissue lung, between diseased tissue breast tumor and healthy tissue breast demonstrates that the human KCNK1 or mRNA can be utilized to diagnose of cancer. Additionally the activity of the human KCNK1 can be modulated to treat cancer.

Inflammatory Diseases

Inflammatory diseases comprise diseases triggered by cellular or non-cellular mediators of the immune system or tissues causing the inflammation of body tissues and subsequently producing an acute or chronic inflammatory condition. Examples for such inflammatory diseases are hypersensitivity reactions of type I – IV, for example but not limited to hypersensitivity diseases of the lung including asthma, atopic diseases, allergic rhinitis or conjunctivitis, angioedema of the

lids, hereditary angioedema, antireceptor hypersensitivity reactions and autoimmune diseases, Hashimoto's thyroiditis, systemic lupus erythematosus, Goodpasture's syndrome, pemphigus, myasthenia gravis, Grave's and Raynaud's disease, type B insulin-resistant diabetes, rheumatoid arthritis, psoriasis, Crohn's disease, scleroderma, mixed connective tissue disease, polymyositis, sarcoidosis, glomerulonephritis, acute or chronic host versus graft reactions.

The human KCNK1 is highly expressed in the following tissues of the immune system and tissues responsive to components of the immune system as well as in the following tissues responsive to mediators of inflammation: bone marrow CD15+ cells. The expression in the above mentioned tissues demonstrates that the human KCNK1 or mRNA can be utilized to diagnose of inflammatory diseases. Additionally the activity of the human KCNK1 can be modulated to treat inflammatory diseases.

Disorders Related to Urology

Genitourinary disorders comprise benign and malign disorders of the organs constituting the genitourinary system of female and male, renal diseases like acute or chronic renal failure, immunologically mediated renal diseases like renal transplant rejection, lupus nephritis, immune complex renal diseases, glomerulopathies, nephritis, toxic nephropathy, obstructive uropathies like benign prostatic hyperplasia (BPH), neurogenic bladder syndrome, urinary incontinence like urge-, stress-, or overflow incontinence, pelvic pain, and erectile dysfunction.

The human KCNK1 is highly expressed in the following urological tissues: spinal cord, prostate, prostate BPH, bladder. The expression in the above mentioned tissues and in particular the differential expression between diseased tissue prostate BPH and healthy tissue prostate demonstrates that the human KCNK1 or mRNA can be utilized to diagnose of urological disorders. Additionally the activity of the human KCNK1 can be modulated to treat urological disorders.

The human KCNK1 is highly expressed in spinal cord tissues: spinal cord. Expression in spinal cord tissues demonstrates that the human KCNK1 or mRNA can be utilized to diagnose of incontinence as an urological disorder. The spinal cord tissues are involved in the neuronal regulation of the urological system. Additionally the activity of the human KCNK1 can be modulated to treat - but not limited to - incontinence.

Applications

The present invention provides for both prophylactic and therapeutic methods for cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases.

The regulatory method of the invention involves contacting a cell with an agent that modulates one or more of the activities of KCNK1. An agent that modulates activity can be an agent as described herein, such as a nucleic acid or a protein, a naturally-occurring cognate ligand of the polypeptide, a peptide, a peptidomimetic, or any small molecule. In one embodiment, the agent stimulates one or more of the biological activities of KCNK1. Examples of such stimulatory agents include the active KCNK1 and nucleic acid molecules encoding a portion of KCNK1. In another embodiment, the agent inhibits one or more of the biological activities of KCNK1. Examples of such inhibitory agents include antisense nucleic acid molecules and antibodies. These regulatory methods can be performed in vitro (e.g., by culturing the cell with the agent) or, alternatively, in vivo (e.g., by administering the agent to a subject). As such, the present invention provides methods of treating an individual afflicted with a disease or disorder characterized by unwanted expression or activity of KCNK1 or a protein in the KCNK1 signaling pathway. In one embodiment, the method involves administering an agent like any agent identified or being identifiable by a screening assay as described herein, or combination of such agents that modulate say upregulate or downregulate the expression or activity of KCNK1 or of any protein in the KCNK1 signaling pathway. In another embodiment, the method involves administering a regulator of KCNK1 as therapy to compensate for reduced or undesirably low expression or activity of KCNK1 or a protein in the KCNK1 signaling pathway.

Stimulation of activity or expression of KCNK1 is desirable in situations in which activity or expression is abnormally low and in which increased activity is likely to have a beneficial effect. Conversely, inhibition of activity or expression of KCNK1 is desirable in situations in which activity or expression of KCNK1 is abnormally high and in which decreasing its activity is likely to have a beneficial effect.

This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents and published patent applications cited throughout this application are hereby incorporated by reference.

Pharmaceutical Compositions

This invention further pertains to novel agents identified by the above-described screening assays and uses thereof for treatments as described herein.

The nucleic acid molecules, polypeptides, and antibodies (also referred to herein as "active compounds") of the invention can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the nucleic acid molecule, protein, or antibody and a pharmaceutically acceptable carrier. As used herein the language "pharma-

aceutically acceptable carrier“ is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

The invention includes pharmaceutical compositions comprising a regulator of KCNK1 expression or activity (and/or a regulator of the activity or expression of a protein in the KCNK1 signaling pathway) as well as methods for preparing such compositions by combining one or more such regulators and a pharmaceutically acceptable carrier. Also within the invention are pharmaceutical compositions comprising a regulator identified using the screening assays of the invention packaged with instructions for use. For regulators that are antagonists of KCNK1 activity or which reduce KCNK1 expression, the instructions would specify use of the pharmaceutical composition for treatment of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases. For regulators that are agonists of KCNK1 activity or increase KCNK1 expression, the instructions would specify use of the pharmaceutical composition for treatment of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases.

An antagonist of KCNK1 may be produced using methods which are generally known in the art. In particular, purified KCNK1 may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind KCNK1. Antibodies to KCNK1 may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies like those which inhibit dimer formation are especially preferred for therapeutic use.

In another embodiment of the invention, the polynucleotides encoding KCNK1, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding KCNK1 may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding KCNK1. Thus, complementary molecules or fragments may be used to modulate KCNK1 activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligonucleotides or larger

fragments can be designed from various locations along the coding or control regions of sequences encoding KCNK1.

Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors which will express nucleic acid sequence complementary to the polynucleotides of the gene encoding KCNK1. These techniques are described, for example, in [Scott and Smith (1990) Science 249:386-390].

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

An additional embodiment of the invention relates to the administration of a pharmaceutical composition containing KCNK1 in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of KCNK1, antibodies to KCNK1, and mimetics, agonists, antagonists, or inhibitors of KCNK1. The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs or hormones.

A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, oral (e.g., inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile

injectable solutions or dispersions. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EMTM (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, a pharmaceutically acceptable polyol like glycerol, propylene glycol, liquid polyethylene glycol, and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin. Sterile injectable solutions can be prepared by incorporating the active compound (e.g., a polypeptide or antibody) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed.

Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or sterotes; a glidant such as

colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

For administration by inhalation, the compounds are delivered in the form of an aerosol spray from a pressurized container or dispenser which contains a suitable propellant, e.g., a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

The compounds can also be prepared in the form of suppositories (e.g., with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the

limitations inherent in the art of compounding such an active compound for the treatment of individuals.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration. For pharmaceutical compositions which include an antagonist of KCNK1 activity, a compound which reduces expression of KCNK1, or a compound which reduces expression or activity of a protein in the KCNK1 signaling pathway or any combination thereof, the instructions for administration will specify use of the composition for cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases. For pharmaceutical compositions which include an agonist of KCNK1 activity, a compound which increases expression of KCNK1, or a compound which increases expression or activity of a protein in the KCNK1 signaling pathway or any combination thereof, the instructions for administration will specify use of the composition for cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases.

15 **Diagnostics**

In another embodiment, antibodies which specifically bind KCNK1 may be used for the diagnosis of disorders characterized by the expression of KCNK1, or in assays to monitor patients being treated with KCNK1 or agonists, antagonists, and inhibitors of KCNK1. Antibodies useful for diagnostic purposes may be prepared in the same manner as those described above for therapeutics. Diagnostic assays for KCNK1 include methods which utilize the antibody and a label to detect KCNK1 in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent joining with a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring KCNK1, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of KCNK1 expression. Normal or standard values for KCNK1 expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, preferably human, with antibody to KCNK1 under conditions suitable for complex formation. The amount of standard complex formation may be quantified by various methods, preferably by photometric means. Quantities of KCNK1 expressed in subject samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding KCNK1 may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantitate gene expression in biopsied tissues in which expression of KCNK1 may be correlated with disease. The diagnostic assay may be used to distinguish between absence, presence, and excess expression of KCNK1, and to monitor regulation of KCNK1 levels during therapeutic intervention.

Polynucleotide sequences encoding KCNK1 may be used for the diagnosis of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases associated with expression of KCNK1. The polynucleotide sequences encoding KCNK1 may be used in Southern, Northern, or dot-blot analysis, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and ELISA assays; and in microarrays utilizing fluids or tissues from patient biopsies to detect altered KCNK1 expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding KCNK1 may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding KCNK1 may be labelled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantitated and compared with a standard value. If the amount of signal in the patient sample is significantly altered from that of a comparable control sample, the nucleotide sequences have hybridized with nucleotide sequences in the sample, and the presence of altered levels of nucleotide sequences encoding KCNK1 in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or in monitoring the treatment of an individual patient.

In order to provide a basis for the diagnosis of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases associated with expression of KCNK1, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding KCNK1, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained from normal samples may be compared with values obtained from

samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Another technique for drug screening which may be used provides for high throughput screening of compounds having suitable binding affinity to the protein of interest as described in published PCT application WO84/03564. In this method, large numbers of different small test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The test compounds are reacted with KCNK1, or fragments thereof, and washed. Bound KCNK1 is then detected by methods well known in the art. Purified KCNK1 can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding KCNK1 specifically compete with a test compound for binding KCNK1. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with KCNK1.

15 **Determination of a Therapeutically Effective Dose**

The determination of a therapeutically effective dose is well within the capability of those skilled in the art. A therapeutically effective dose refers to that amount of active ingredient which increases or decreases KCNK1 activity relative to KCNK1 activity which occurs in the absence of the therapeutically effective dose. For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays or in animal models, usually mice, rabbits, dogs, or pigs. The animal model also can be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

Therapeutic efficacy and toxicity, e.g., ED_{50} (the dose therapeutically effective in 50% of the population) and LD_{50} (the dose lethal to 50% of the population), can be determined by standard pharmaceutical procedures in cell cultures or experimental animals. The dose ratio of toxic to therapeutic effects is the therapeutic index, and it can be expressed as the ratio, LD_{50}/ED_{50} . Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that include the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration. The exact dosage will be determined by the practitioner, in light of factors related

to the subject that requires treatment. Dosage and administration are adjusted to provide sufficient levels of the active ingredient or to maintain the desired effect. Factors which can be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long-acting pharmaceutical compositions can be administered every 3 to 4 days, every week, or once every two weeks depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts can vary from 0.1 micrograms to 100,000 micrograms, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc. If the reagent is a single-chain antibody, polynucleotides encoding the antibody can be constructed and introduced into a cell either *ex vivo* or *in vivo* using well-established techniques including, but not limited to, transferrin-polycation-mediated DNA transfer, transfection with naked or encapsulated nucleic acids, liposome-mediated cellular fusion, intracellular transportation of DNA-coated latex beads, protoplast fusion, viral infection, electroporation, "gene gun", and DEAE- or calcium phosphate-mediated transfection.

If the expression product is mRNA, the reagent is preferably an antisense oligonucleotide or a ribozyme. Polynucleotides which express antisense oligonucleotides or ribozymes can be introduced into cells by a variety of methods, as described above. Preferably, a reagent reduces expression of KCNK1 gene or the activity of KCNK1 by at least about 10, preferably about 50, more preferably about 75, 90, or 100% relative to the absence of the reagent. The effectiveness of the mechanism chosen to decrease the level of expression of KCNK1 gene or the activity of KCNK1 can be assessed using methods well known in the art, such as hybridization of nucleotide probes to KCNK1-specific mRNA, quantitative RT-PCR, immunologic detection of KCNK1, or measurement of KCNK1 activity.

In any of the embodiments described above, any of the pharmaceutical compositions of the invention can be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy can be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents can act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects. Any of the therapeutic

methods described above can be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

5 Nucleic acid molecules of the invention are those nucleic acid molecules which are contained in a group of nucleic acid molecules consisting of (i) nucleic acid molecules encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 2, (ii) nucleic acid molecules comprising the sequence of SEQ ID NO: 1, (iii) nucleic acid molecules having the sequence of SEQ ID NO: 1, (iv) nucleic acid molecules the complementary strand of which hybridizes under stringent conditions to a nucleic acid molecule of (i), (ii), or (iii); and (v) nucleic acid molecules the
10 sequence of which differs from the sequence of a nucleic acid molecule of (iii) due to the degeneracy of the genetic code, wherein the polypeptide encoded by said nucleic acid molecule has KCNK1 activity.

Polypeptides of the invention are those polypeptides which are contained in a group of polypeptides consisting of (i) polypeptides having the sequence of SEQ ID NO: 2, (ii) polypeptides
15 comprising the sequence of SEQ ID NO: 2, (iii) polypeptides encoded by nucleic acid molecules of the invention and (iv) polypeptides which show at least 99%, 98%, 95%, 90%, or 80% homology with a polypeptide of (i), (ii), or (iii), wherein said purified polypeptide has KCNK1 activity.

An object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological
20 diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising the steps of (i) contacting a test compound with a KCNK1 polypeptide, (ii) detect binding of said test compound to said KCNK1 polypeptide. E.g., compounds that bind to the KCNK1 polypeptide are identified potential therapeutic agents for such a disease.

25 Another object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising the steps of (i) determining the activity of a KCNK1 polypeptide at a certain concentration of a test compound or
30 in the absence of said test compound, (ii) determining the activity of said polypeptide at a different concentration of said test compound. E.g., compounds that lead to a difference in the activity of the KCNK1 polypeptide in (i) and (ii) are identified potential therapeutic agents for such a disease.

Another object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising the steps of (i) determining the activity of a KCNK1 polypeptide at a certain concentration of a test compound, (ii) determining the activity of a KCNK1 polypeptide at the presence of a compound known to be a regulator of a KCNK1 polypeptide. E.g., compounds that show similar effects on the activity of the KCNK1 polypeptide in (i) as compared to compounds used in (ii) are identified potential therapeutic agents for such a disease.

Another object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising i) cell expressing KCNK1 having functional activity, ii) contacting said cell with the test compounds, iii) measuring functional activity.

Another object of the invention is the method of the above, where the cell is expressing endogenous KCNK1.

Another object of the invention are the methods of the above, where the cell is expressing recombinant KCNK1.

Another object of the invention are the methods of the above, where the measured functional activity is the KCNK1 current.

Another object of the invention are the methods of the above, where the functional activity is a change in membrane potential.

Another object of the invention are the methods of the above, where the functional activity is a change in ion concentration.

Other objects of the invention are methods of the above, wherein the step of contacting is in or at the surface of a cell.

Other objects of the invention are methods of the above, wherein the cell is in vitro.

Other objects of the invention are methods of the above, wherein the step of contacting is in a cell-free system.

Other objects of the invention are methods of the above, wherein the polypeptide is coupled to a detectable label.

Other objects of the invention are methods of the above, wherein the compound is coupled to a detectable label.

- 5 Other objects of the invention are methods of the above, wherein the test compound displaces a ligand which is first bound to the polypeptide. Preferably the ligand to be displaced is coupled to a detectable label.

Other objects of the invention are methods of the above, wherein the polypeptide is attached to a solid support.

- 10 Other objects of the invention are methods of the above, wherein the compound is attached to a solid support.

- Another object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising the steps of (i) contacting a test compound with a KCNK1 polynucleotide, (ii) detect binding of said test compound to said KCNK1 polynucleotide. Compounds that, e.g., bind to the KCNK1 polynucleotide are potential therapeutic agents for the treatment of such diseases.
- 15

- Another object of the invention is the method of the above, wherein the nucleic acid molecule is RNA.
- 20

Another object of the invention is a method of the above, wherein the contacting step is in or at the surface of a cell.

Another object of the invention is a method of the above, wherein the contacting step is in a cell-free system.

- 25 Another object of the invention is a method of the above, wherein the polynucleotide is coupled to a detectable label.

Another object of the invention is a method of the above, wherein the test compound is coupled to a detectable label.

Another object of the invention is a method of diagnosing a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising the steps of (i) determining the amount of a KCNK1 polynucleotide in a sample taken from said mammal, (ii) determining the amount of KCNK1 polynucleotide in healthy and/or diseased mammal. A disease is diagnosed, e.g., if there is a substantial similarity in the amount of KCNK1 polynucleotide in said test mammal as compared to a diseased mammal.

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising a therapeutic agent which binds to a KCNK1 polypeptide.

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising a therapeutic agent which regulates the activity of a KCNK1 polypeptide.

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising a therapeutic agent which regulates the activity of a KCNK1 polypeptide, wherein said therapeutic agent is (i) a small molecule, (ii) an RNA molecule, (iii) an antisense oligonucleotide, (iv) a polypeptide, (v) an antibody, or (vi) a ribozyme.

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising a KCNK1 polynucleotide.

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising a KCNK1 polypeptide.

Another object of the invention is the use of regulators of a KCNK1 for the preparation of a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal.

Another object of the invention is a method for the preparation of a pharmaceutical composition useful for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising the steps of (i) identifying a regulator of KCNK1, (ii) determining whether said regulator ameliorates the symptoms of a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal; and (iii) combining of said regulator with an acceptable pharmaceutical carrier.

Another object of the invention is the use of a regulator of KCNK1 for the regulation of KCNK1 activity in a mammal having a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases.

The expression of human KCNK1 in cardiovascular and cancer related tissues (as described above) suggests a particular – but not limited to – utilization of KCNK1 for diagnosis and modulation of cardiovascular diseases and cancer. Furthermore the above described expression suggest a – but not limited to – utilization of KCNK1 to dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, inflammation and urological diseases.

The examples below are provided to illustrate the subject invention. These examples are provided by way of illustration and are not included for the purpose of limiting the invention.

Examples**Example 1: Search for homologous sequences in public sequence data bases**

The degree of homology can readily be calculated by known methods. Preferred methods to determine homology are designed to give the largest match between the sequences tested.

- 5 Methods to determine homology are codified in publicly available computer programs such as BestFit, BLASTP, BLASTN, and FASTA. The BLAST programs are publicly available from NCBI and other sources in the internet.

- For KCNK1 the following hits to known sequences were identified by using the BLAST algorithm [Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ; Nucleic Acids
10 Res 1997 Sep 1; 25(17): 3389-402] and the following set of parameters: matrix = BLOSUM62 and low complexity filter. The following databases were searched: NCBI (non-redundant database) and DERWENT patent database (Geneseq).

The following hits were found:

- >ref[NM_002245.2] Homo sapiens potassium channel, subfamily K, member 1 (KCNK1), mRNA
15 Length = 1901, Score = 3350 bits (1742), Expect = 0.0, Identities = 1770/1798 (98%)
- >NA2002:ABL63324 Abl63324 Breast cancer related gene sequence SEQ IDNO:1661. 5/2002
Length = 1882, Score = 3313 bits (1723), Expect = 0.0, Identities = 1751/1779 (98%)
- >NA2002:ABL63116 Abl63116 Breast cancer related gene sequence SEQ IDNO:1453. 5/2002
Length = 1882, Score = 3313 bits (1723), Expect = 0.0, Identities = 1751/1779 (98%)
- 20 >NA2002:ABL62908 Abl62908 Breast cancer related gene sequence SEQ IDNO:1245. 5/2002
Length = 1882, Score = 3313 bits (1723), Expect = 0.0, Identities = 1751/1779 (98%)
- >NA2000:AAA27745 Aaa27745 Human potassium channel TWIK-1 cDNA. 8/2000
Length = 1894, Score = 3313 bits (1723), Expect = 0.0, Identities = 1751/1779 (98%)
- >NA1997:AAT64960 Aat64960 TWIK-1 potassium channel cDNA. 3/1998
25 Length = 1894, Score = 3313 bits (1723), Expect = 0.0, Identities = 1751/1779 (98%)
- >gb[AR175707.1]AR175707 Sequence 1 from patent US 6309855
Length = 1894, Score = 3313 bits (1723), Expect = 0.0, Identities = 1751/1779 (98%)

>emb|AX331152.1| Sequence 1661 from Patent WO0194629

Length = 1882, Score = 3313 bits (1723), Expect = 0.0, Identities = 1751/1779 (98%)

>emb|AX330944.1| Sequence 1453 from Patent WO0194629

Length = 1882, Score = 3313 bits (1723), Expect = 0.0, Identities = 1751/1779 (98%)

5 >emb|AX330736.1| Sequence 1245 from Patent WO0194629

Length = 1882, Score = 3313 bits (1723), Expect = 0.0, Identities = 1751/1779 (98%).

Example 2: Expression profiling

Total cellular RNA was isolated from cells by one of two standard methods: 1) guanidine isothiocyanate/Cesium chloride density gradient centrifugation [Kellogg, (1990)] ; or with the Tri-
10 Reagent protocol according to the manufacturer's specifications (Molecular Research Center, Inc., Cincinnati, Ohio). Total RNA prepared by the Tri-reagent protocol was treated with DNase I to remove genomic DNA contamination.

For relative quantitation of the mRNA distribution of KCNK1, total RNA from each cell or tissue source was first reverse transcribed. 85 µg of total RNA was reverse transcribed using 1 µmole
15 random hexamer primers, 0.5 mM each of dATP, dCTP, dGTP and dTTP (Qiagen, Hilden, Germany), 3000 U RnaseQut (Invitrogen, Groningen, Netherlands) in a final volume of 680 µl. The first strand synthesis buffer and Omniscript reverse transcriptase (2 u/µl) were from (Qiagen, Hilden, Germany). The reaction was incubated at 37°C for 90 minutes and cooled on ice. The volume was adjusted to 6800 µl with water, yielding a final concentration of 12.5 ng/µl of starting
20 RNA.

For relative quantitation of the distribution of KCNK1 mRNA in cells and tissues the Applied Biosystems 7900 HT Sequence Detection system or Biorad iCycler was used according to the manufacturer's specifications and protocols. PCR reactions were set up to quantitate KCNK1 and the housekeeping genes HPRT (hypoxanthine phosphoribosyltransferase), GAPDH (glyceraldehyde-3-phosphate dehydrogenase), β-actin, and others. Forward and reverse primers and probes for KCNK1
25 were designed using the Perkin Elmer ABI Primer Express™ software and were synthesized by TibMolBiol (Berlin, Germany). The KCNK1 forward primer sequence was: Primer1 (SEQ ID NO: 3). The KCNK1 reverse primer sequence was Primer2 (SEQ ID NO: 4). Probe1 (SEQ ID NO: 5), labelled with FAM (carboxyfluorescein succinimidyl ester) as the reporter dye and TAMRA (carboxytetramethylrhodamine) as the quencher, is used as a probe for KCNK1. The following
30 reagents were prepared in a total of 25 µl : 1x TaqMan buffer A, 5.5 mM MgCl₂, 200 nM of dATP, dCTP, dGTP, and dUTP, 0.025 U/µl AmpliTaq Gold™, 0.01 U/ µl AmpErase and Probe1 (SEQ ID

NO: 4), KCNK1 forward and reverse primers each at 200 nM, 200 nM KCNK1 FAM/TAMRA-labelled probe, and 5 µl of template cDNA. Thermal cycling parameters were 2 min at 50°C, followed by 10 min at 95°C, followed by 40 cycles of melting at 95°C for 15 sec and annealing/extending at 60°C for 1 min.

5 *Calculation of corrected CT values*

The CT (threshold cycle) value is calculated as described in the “Quantitative determination of nucleic acids” section. The CF-value (factor for threshold cycle correction) is calculated as follows :

1. PCR reactions were set up to quantitate the housekeeping genes (HKG) for each cDNA sample.
2. CT_{HKG} -values (threshold cycle for housekeeping gene) were calculated as described in the “Quantitative determination of nucleic acids” section.
3. CT_{HKG} -mean values (CT mean value of all HKG tested on one cDNAs) of all HKG for each cDNA are calculated (n = number of HKG):

$$CT_{HKG-n}\text{-mean value} = (CT_{HKG1}\text{-value} + CT_{HKG2}\text{-value} + \dots + CT_{HKG-n}\text{-value}) / n$$
4. CT_{panel} mean value (CT mean value of all HKG in all tested cDNAs) =

$$(CT_{HKG1}\text{-mean value} + CT_{HKG2}\text{-mean value} + \dots + CT_{HKG-y}\text{-mean value}) / y$$
(y = number of cDNAs)
5. CF_{cDNA-n} (correction factor for cDNA n) = $CT_{\text{panel}}\text{-mean value} - CT_{HKG-n}\text{-mean value}$
6. CT_{cDNA-n} (CT value of the tested gene for the cDNA n) + CF_{cDNA-n} (correction factor for cDNA n) = $CT_{\text{cor-cDNA-n}}$ (corrected CT value for a gene on cDNA n)

Calculation of relative expression

Definition : highest $CT_{\text{cor-cDNA-n}} \neq 40$ is defined as $CT_{\text{cor-cDNA}} [\text{high}]$

Relative Expression = $2^{(CT_{\text{cor-cDNA}} [\text{high}] - CT_{\text{cor-cDNA-n}})}$

25 *Tissues*

The expression of KCNK1 was investigated in the tissues listed in table 1.

Expression profile

The results of the the mRNA-quantification (expression profiling) is shown in Table 1.

Table 1: Relative expression of KCNK1 in various human tissues

	fetal heart	52
	heart	146
5	pericardium	76
	heart atrium (right)	838
	heart atrium (left)	1531
	heart ventricle (left)	798
	interventricular septum	208
10	fetal aorta	7
	aorta	147
	artery	30
	coronary artery	1
	vein	33
15	coronary artery smooth muscle primary cells	43
	HUVEC cells	7
	skin	168
20	adrenal gland	40
	thyroid	124
	thyroid tumor	241
	pancreas	125
	pancreas liver cirrhosis	115
25	esophagus	431
	esophagus tumor	45
	stomach	108
	stomach tumor	127
30	colon	413
	colon tumor	260
	small intestine	127
	ileum	286
	ileum tumor	2
35	ileum chronic inflammation	2
	rectum	108
	salivary gland	176
	fetal liver	16

	liver	48
	liver liver cirrhosis	129
	liver tumor	6
	HEP G2 cells	2
5		
	leukocytes (peripheral blood)	1
	Jurkat (T-cells)	63
	bone marrow	2
	erythrocytes	6
10	lymphnode	21
	thymus	7
	thrombocytes	65
	bone marrow CD71+ cells	49
	bone marrow CD34+ cells	15
15	bone marrow CD15+ cells	125
	cord blood CD71+ cells	133
	spleen	8
	spleen liver cirrhosis	15
20		
	skeletal muscle	13
	adipose	79
	fetal brain	92
	brain	809
25	Alzheimer brain	630
	cerebellum	13873
	cerebellum (right)	4905
	cerebellum (left)	2856
	cerebral cortex	1898
30	Alzheimer cerebral cortex	4939
	frontal lobe	1531
	Alzheimer brain frontal lobe	534
	occipital lobe	3236
	parietal lobe	2574
35	temporal lobe	3019
	precentral gyrus	2702
	postcentral gyrus	3717
	tonsilla cerebelli	5185
	vermis cerebelli	3421

	pons	425
	substantia nigra	4
	cerebral meninges	49
	cerebral peduncles	6517
5	corpus callosum	729
	hippocampus	3373
	thalamus	326
	dorsal root ganglia	25
	spinal cord	251
10	neuroblastoma SK-N-MC cells	4
	neuroblastoma SH-SY5Y cells	122
	neuroblastoma IMR32 cells	537
	retina	12
15	fetal lung	190
	fetal lung fibroblast IMR-90 cells	2
	lung	32
	lung tumor	278
	lung COPD	15
20	trachea	284
	cervix	218
	testis	26
	HeLa cells (cervix tumor)	87
25	placenta	68
	uterus	65
	uterus tumor	75
	ovary tumor	83
	breast	1053
30	breast tumor	59
	MDA MB 231 cells (breast tumor)	56
	mammary gland	49
	prostate	185
35	prostate BPH	4
	bladder	122
	penis	21
	corpus cavernosum	29
	fetal kidney	114

kidney	73
kidney tumor	12
HEK 293 cells	202

Example 3: Antisense Analysis

5 Knowledge of the correct, complete cDNA sequence coding for KCNK1 enables its use as a tool for antisense technology in the investigation of gene function. Oligonucleotides, cDNA or genomic fragments comprising the antisense strand of a polynucleotide coding for KCNK1 are used either in vitro or in vivo to inhibit translation of the mRNA. Such technology is now well known in the art, and antisense molecules can be designed at various locations along the nucleotide sequences. By treatment of cells or whole test animals with such antisense sequences, the gene of interest is effectively turned off. Frequently, the function of the gene is ascertained by observing behavior at the intracellular, cellular, tissue or organismal level (e.g., lethality, loss of differentiated function, changes in morphology, etc.).

15 In addition to using sequences constructed to interrupt transcription of a particular open reading frame, modifications of gene expression is obtained by designing antisense sequences to intron regions, promoter/enhancer elements, or even to trans-acting regulatory genes.

Example 4: Expression of KCNK1

Expression of KCNK1 is accomplished by subcloning the cDNAs into appropriate expression vectors and transfecting the vectors into expression hosts such as, e.g., *E. coli*. In a particular case, the vector is engineered such that it contains a promoter for β -galactosidase, upstream of the cloning site, followed by sequence containing the amino-terminal Methionine and the subsequent seven residues of β -galactosidase. Immediately following these eight residues is an engineered bacteriophage promoter useful for artificial priming and transcription and for providing a number of unique endonuclease restriction sites for cloning.

25 Induction of the isolated, transfected bacterial strain with Isopropyl- β -D-thiogalactopyranoside (IPTG) using standard methods produces a fusion protein corresponding to the first seven residues of β -galactosidase, about 15 residues of "linker", and the peptide encoded within the cDNA. Since cDNA clone inserts are generated by an essentially random process, there is probability of 33% that the included cDNA will lie in the correct reading frame for proper translation. If the cDNA is not in the proper reading frame, it is obtained by deletion or insertion of the appropriate number of bases using well known methods including in vitro mutagenesis, digestion with exonuclease III or mung bean nuclease, or the inclusion of an oligonucleotide linker of appropriate length.

The KCNK1 cDNA is shuttled into other vectors known to be useful for expression of proteins in specific hosts. Oligonucleotide primers containing cloning sites as well as a segment of DNA (about 25 bases) sufficient to hybridize to stretches at both ends of the target cDNA is synthesized chemically by standard methods. These primers are then used to amplify the desired gene segment by PCR. The resulting gene segment is digested with appropriate restriction enzymes under standard conditions and isolated by gel electrophoresis. Alternately, similar gene segments are produced by digestion of the cDNA with appropriate restriction enzymes. Using appropriate primers, segments of coding sequence from more than one gene are ligated together and cloned in appropriate vectors. It is possible to optimize expression by construction of such chimeric sequences.

Suitable expression hosts for such chimeric molecules include, but are not limited to, mammalian cells such as Chinese Hamster Ovary (CHO) and human 293 cells., insect cells such as Sf9 cells, yeast cells such as *Saccharomyces cerevisiae* and bacterial cells such as *E. coli*. For each of these cell systems, a useful expression vector also includes an origin of replication to allow propagation in bacteria, and a selectable marker such as the β -lactamase antibiotic resistance gene to allow plasmid selection in bacteria. In addition, the vector may include a second selectable marker such as the neomycin phosphotransferase gene to allow selection in transfected eukaryotic host cells. Vectors for use in eukaryotic expression hosts require RNA processing elements such as 3' polyadenylation sequences if such are not part of the cDNA of interest.

Additionally, the vector contains promoters or enhancers which increase gene expression. Such promoters are host specific and include MMTV, SV40, and metallothionine promoters for CHO cells; trp, lac, tac and T7 promoters for bacterial hosts; and alpha factor, alcohol oxidase and PGH promoters for yeast. Transcription enhancers, such as the rous sarcoma virus enhancer, are used in mammalian host cells. Once homogeneous cultures of recombinant cells are obtained through standard culture methods, large quantities of recombinantly produced KCNK1 are recovered from the conditioned medium and analyzed using chromatographic methods known in the art. For example, KCNK1 can be cloned into the expression vector pcDNA3, as exemplified herein. This product can be used to transform, for example, HEK293 or COS by methodology standard in the art. Specifically, for example, using Lipofectamine (Gibco BRL catalog no. 18324-020) mediated gene transfer.

Example 5: Isolation of Recombinant KCNK1

KCNK1 is expressed as a chimeric protein with one or more additional polypeptide domains added to facilitate protein purification. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on

immobilized metals [Appa Rao, 1997] and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Washington). The inclusion of a cleavable linker sequence such as Factor Xa or enterokinase (Invitrogen, Groningen, The Netherlands) between the purification domain and the KCNK1 sequence is useful to facilitate expression of KCNK1.

5 **Example 6: Production of KCNK1 Specific Antibodies**

Two approaches are utilized to raise antibodies to KCNK1, and each approach is useful for generating either polyclonal or monoclonal antibodies. In one approach, denatured protein from reverse phase HPLC separation is obtained in quantities up to 75 mg. This denatured protein is used to immunize mice or rabbits using standard protocols; about 100 µg are adequate for
10 immunization of a mouse, while up to 1 mg might be used to immunize a rabbit. For identifying mouse hybridomas, the denatured protein is radioiodinated and used to screen potential murine B-cell hybridomas for those which produce antibody. This procedure requires only small quantities of protein, such that 20 mg is sufficient for labeling and screening of several thousand clones.

In the second approach, the amino acid sequence of an appropriate KCNK1 domain, as deduced
15 from translation of the cDNA, is analyzed to determine regions of high antigenicity. Oligopeptides comprising appropriate hydrophilic regions are synthesized and used in suitable immunization protocols to raise antibodies. The optimal amino acid sequences for immunization are usually at the C-terminus, the N-terminus and those intervening, hydrophilic regions of the polypeptide which are likely to be exposed to the external environment when the protein is in its natural
20 conformation.

Typically, selected peptides, about 15 residues in length, are synthesized using an Applied Biosystems Peptide Synthesizer Model 431A using fmoc-chemistry and coupled to keyhole limpet hemocyanin (KLH; Sigma, St. Louis, MO) by reaction with M-maleimidobenzoyl-N-hydroxy-succinimide ester, MBS. If necessary, a cysteine is introduced at the N-terminus of the peptide to
25 permit coupling to KLH. Rabbits are immunized with the peptide-KLH complex in complete Freund's adjuvant. The resulting antisera are tested for antipeptide activity by binding the peptide to plastic, blocking with 1% bovine serum albumin, reacting with antisera, washing and reacting with labeled (radioactive or fluorescent), affinity purified, specific goat anti-rabbit IgG.

Hybridomas are prepared and screened using standard techniques. Hybridomas of interest are
30 detected by screening with labeled KCNK1 to identify those fusions producing the monoclonal antibody with the desired specificity. In a typical protocol, wells of plates (FAST; Becton-Dickinson, Palo Alto, CA) are coated during incubation with affinity purified, specific rabbit anti-mouse (or suitable antispecies 1 g) antibodies at 10 mg/ml. The coated wells are blocked with 1%

bovine serum albumin, (BSA), washed and incubated with supernatants from hybridomas. After washing the wells are incubated with labeled KCNK1 at 1 mg/ml. Supernatants with specific antibodies bind more labeled KCNK1 than is detectable in the background. Then clones producing specific antibodies are expanded and subjected to two cycles of cloning at limiting
5 dilution. Cloned hybridomas are injected into pristane-treated mice to produce ascites, and monoclonal antibody is purified from mouse ascitic fluid by affinity chromatography on Protein A. Monoclonal antibodies with affinities of at least

10^8 M^{-1} , preferably 10^9 to 10^{10} M^{-1} or stronger, are typically made by standard procedures.

Example 7: Diagnostic Test Using KCNK1 Specific Antibodies

10 Particular KCNK1 antibodies are useful for investigating signal transduction and the diagnosis of infectious or hereditary conditions which are characterized by differences in the amount or distribution of KCNK1 or downstream products of an active signaling cascade.

Diagnostic tests for KCNK1 include methods utilizing antibody and a label to detect KCNK1 in human body fluids, membranes, cells, tissues or extracts of such. The polypeptides and antibodies
15 of the present invention are used with or without modification. Frequently, the polypeptides and antibodies are labeled by joining them, either covalently or noncovalently, with a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known and have been reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent agents, chemi-
20 luminescent agents, chromogenic agents, magnetic particles and the like.

A variety of protocols for measuring soluble or membrane-bound KCNK1, using either polyclonal or monoclonal antibodies specific for the protein, are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA) and fluorescent activated cell sorting (FACS). A two-site monoclonal-based immunoassay utilizing monoclonal antibodies
25 reactive to two non-interfering epitopes on KCNK1 is preferred, but a competitive binding assay may be employed.

Example 8: Purification of Native KCNK1 Using Specific Antibodies

Native or recombinant KCNK1 is purified by immunoaffinity chromatography using antibodies specific for KCNK1. In general, an immunoaffinity column is constructed by covalently coupling the anti-TRH antibody to an activated chromatographic resin.

- 5 Polyclonal immunoglobulins are prepared from immune sera either by precipitation with ammonium sulfate or by purification on immobilized Protein A (Pharmacia LKB Biotechnology, Piscataway N.J.). Likewise, monoclonal antibodies are prepared from mouse ascites fluid by ammonium sulfate precipitation or chromatography on immobilized Protein A. Partially purified immunoglobulin is covalently attached to a chromatographic resin such as CnBr-activated
10 Sepharose (Pharmacia LKB Biotechnology). The antibody is coupled to the resin, the resin is blocked, and the derivative resin is washed according to the manufacturer's instructions.

- Such immunoaffinity columns are utilized in the purification of KCNK1 by preparing a fraction from cells containing KCNK1 in a soluble form. This preparation is derived by solubilization of whole cells or of a subcellular fraction obtained via differential centrifugation (with or without
15 addition of detergent) or by other methods well known in the art. Alternatively, soluble KCNK1 containing a signal sequence is secreted in useful quantity into the medium in which the cells are grown.

- A soluble KCNK1-containing preparation is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of KCNK1 (e.g., high
20 ionic strength buffers in the presence of detergent). Then, the column is eluted under conditions that disrupt antibody/protein binding (e.g., a buffer of pH 2-3 or a high concentration of a chaotrope such as urea or thiocyanate ion), and KCNK1 is collected.

Example 9: Drug Screening

- This invention is particularly useful for screening therapeutic compounds by using KCNK1 or
25 binding fragments thereof in any of a variety of drug screening techniques. As KCNK1 is an electrogenic target any of the methods commonly used in the art may potentially be used to identify KCNK1 ligands modulating functional activity of the channel. For example, the functional activity of ion channels such as KCNK1 can be measured using any of a variety of appropriate functional assays in which modulation of the channel results in an observable change
30 in membrane potential, ion current, ion concentration across the membrane, or any other measurable change exerted by KCNK1. Alternatively, the polypeptide or fragment can be employed in a binding assay either free in solution, affixed to a solid support or expressed on a cell

surface. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the polypeptide or fragment. Drugs are screened against such transformed cells in competitive binding assays. Such cells, either in viable or fixed form, are used for standard binding assays.

- 5 Measured, for example, is the formation of complexes between KCNK1 and the agent being tested. Alternatively, one examines the diminution in complex formation between KCNK1 and a labelled ligand caused by the agent being tested.

Thus, the present invention provides methods of screening for drug candidates, drugs, or any other agents which affects functional activity by binding to the channel protein. These methods, well
10 known in the art, comprise contacting such an agent with KCNK1 polypeptide or a fragment thereof and assaying (i) for the presence of a complex between the agent and KCNK1 polypeptide or fragment, or (ii) for the presence of a complex between KCNK1 polypeptide or fragment and the cell. In such competitive binding assays, the KCNK1 polypeptide or fragment is typically labeled. After suitable incubation, free KCNK1 polypeptide or fragment is separated from that
15 present in bound form, and the amount of free or uncomplexed label is a measure of the ability of the particular agent to bind to KCNK1 or to interfere with the KCNK1-agent complex.

Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to KCNK1 polypeptides. Briefly stated, large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other
20 surface. The peptide test compounds are reacted with KCNK1 polypeptide and washed. Bound KCNK1 polypeptide is then detected by methods well known in the art. Purified KCNK1 are also coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies are used to capture the peptide and immobilize it on the solid support.

This invention also contemplates the use of competitive drug screening assays in which
25 neutralizing antibodies capable of binding KCNK1 specifically compete with a test compound for binding to KCNK1 polypeptides or fragments thereof. In this manner, the antibodies are used to detect the presence of any peptide which shares one or more antigenic determinants with KCNK1.

Example 10: Rational Drug Design

The goal of rational drug design is to produce structural analogs of biologically active polypeptides of interest or of small molecules with which they interact, agonists, antagonists, or inhibitors. Any of these examples are used to fashion drugs which are more active or stable forms of the polypeptide or which enhance or interfere with the function of a polypeptide in vivo.

In one approach, the three-dimensional structure of a protein of interest, or of a protein-inhibitor complex, is determined by x-ray crystallography, by computer modeling or, most typically, by a combination of the two approaches. Both the shape and charges of the polypeptide must be ascertained to elucidate the structure and to determine active site(s) of the molecule. Less often, useful information regarding the structure of a polypeptide is gained by modeling based on the structure of homologous proteins. In both cases, relevant structural information is used to design efficient inhibitors. Useful examples of rational drug design include molecules which have improved activity or stability or which act as inhibitors, agonists, or antagonists of native peptides.

It is also possible to isolate a target-specific antibody, selected by functional assay, as described above, and then to solve its crystal structure. This approach, in principle, yields a pharmacore upon which subsequent drug design is based. It is possible to bypass protein crystallography altogether by generating anti-idiotypic antibodies (anti-ids) to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site of the anti-ids is expected to be an analog of the original receptor. The anti-id is then used to identify and isolate peptides from banks of chemically or biologically produced peptides. The isolated peptides then act as the pharmacore.

By virtue of the present invention, sufficient amount of polypeptide are made available to perform such analytical studies as X-ray crystallography. In addition, knowledge of the KCNK1 amino acid sequence provided herein provides guidance to those employing computer modeling techniques in place of or in addition to x-ray crystallography.

Example 11: Use and Administration of Antibodies, Inhibitors, or Antagonists

Antibodies, inhibitors, or antagonists of KCNK1 or other treatments and compounds that are limiters of signal transduction (LSTs), provide different effects when administered therapeutically. LSTs are formulated in a nontoxic, inert, pharmaceutically acceptable aqueous carrier medium preferably at a pH of about 5 to 8, more preferably 6 to 8, although pH may vary according to the characteristics of the antibody, inhibitor, or antagonist being formulated and the condition to be treated. Characteristics of LSTs include solubility of the molecule, its half-life and antigenicity/-

immunogenicity. These and other characteristics aid in defining an effective carrier. Native human proteins are preferred as LSTs, but organic or synthetic molecules resulting from drug screens are equally effective in particular situations.

5 LSTs are delivered by known routes of administration including but not limited to topical creams and gels; transmucosal spray and aerosol; transdermal patch and bandage; injectable, intravenous and lavage formulations; and orally administered liquids and pills particularly formulated to resist stomach acid and enzymes. The particular formulation, exact dosage, and route of administration is determined by the attending physician and varies according to each specific situation.

10 Such determinations are made by considering multiple variables such as the condition to be treated, the LST to be administered, and the pharmacokinetic profile of a particular LST. Additional factors which are taken into account include severity of the disease state, patient's age, weight, gender and diet, time and frequency of LST administration, possible combination with other drugs, reaction sensitivities, and tolerance/response to therapy. Long acting LST formulations might be administered every 3 to 4 days, every week, or once every two weeks
15 depending on half-life and clearance rate of the particular LST.

Normal dosage amounts vary from 0.1 to 10^5 μg , up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature; see U.S. Pat. Nos. 4,657,760; 5,206,344; or 5,225,212. Those skilled in the art employ different formulations for different LSTs. Administration to cells such as nerve cells
20 necessitates delivery in a manner different from that to other cells such as vascular endothelial cells.

It is contemplated that abnormal signal transduction, trauma, or diseases which trigger KCNK1 activity are treatable with LSTs. These conditions or diseases are specifically diagnosed by the tests discussed above, and such testing should be performed in suspected cases of viral, bacterial or
25 fungal infections, allergic responses, mechanical injury associated with trauma, hereditary diseases, lymphoma or carcinoma, or other conditions which activate the genes of lymphoid or neuronal tissues.

Example 12: Production of Non-human Transgenic Animals

Animal model systems which elucidate the physiological and behavioral roles of the KCNK1 are
30 produced by creating nonhuman transgenic animals in which the activity of the KCNK1 is either increased or decreased, or the amino acid sequence of the expressed KCNK1 is altered, by a variety of techniques. Examples of these techniques include, but are not limited to: 1) Insertion of

normal or mutant versions of DNA encoding a KCNK1, by microinjection, electroporation, retroviral transfection or other means well known to those skilled in the art, into appropriately fertilized embryos in order to produce a transgenic animal or 2) homologous recombination of mutant or normal, human or animal versions of these genes with the native gene locus in transgenic animals to alter the regulation of expression or the structure of these KCNK1 sequences. The technique of homologous recombination is well known in the art. It replaces the native gene with the inserted gene and hence is useful for producing an animal that cannot express native KCNK1s but does express, for example, an inserted mutant KCNK1, which has replaced the native KCNK1 in the animal's genome by recombination, resulting in underexpression of the transporter. Microinjection adds genes to the genome, but does not remove them, and the technique is useful for producing an animal which expresses its own and added KCNK1, resulting in overexpression of the KCNK1.

One means available for producing a transgenic animal, with a mouse as an example, is as follows: Female mice are mated, and the resulting fertilized eggs are dissected out of their oviducts. The eggs are stored in an appropriate medium such as cesiumchloride M2 medium. DNA or cDNA encoding KCNK1 is purified from a vector by methods well known to the one skilled in the art. Inducible promoters may be fused with the coding region of the DNA to provide an experimental means to regulate expression of the transgene. Alternatively or in addition, tissue specific regulatory elements may be fused with the coding region to permit tissue-specific expression of the transgene. The DNA, in an appropriately buffered solution, is put into a microinjection needle (which may be made from capillary tubing using a piper puller) and the egg to be injected is put in a depression slide. The needle is inserted into the pronucleus of the egg, and the DNA solution is injected. The injected egg is then transferred into the oviduct of a pseudopregnant mouse which is a mouse stimulated by the appropriate hormones in order to maintain false pregnancy, where it proceeds to the uterus, implants, and develops to term. As noted above, microinjection is not the only method for inserting DNA into the egg but is used here only for exemplary purposes.

References

- U.S. 5,641,673
- U.S. 5,283,317
- 30 U.S. 5,565,332
- US 6309855
- WO 92/01810
- WO 94/13804
- WO 01/94629

- Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ; *Nucleic Acids Res* 1997 Sep 1; 25(17): 3389-402
- Barnes, 2000, *Chest*, 117:10S14S
- Berman et al. 2000: *J. Neurochem.* 74 (4), 1443-1451
- 5 Baxter et al. 2002: *J. Biomol. Screen.* 7(1): 79-85
- Colbere-Garapin *et al.*, 1981, *J. Mol. Biol.* 150, 1-14
- Gergen and Weiss, 1992, *Am Rev Respir Dis* 146:823-824
- Gibson et al., 1996, *Genome Research* 6: 995-1001
- Hahnenberger et al., *Nature Biotech.* 14, 880-884
- 10 Haseloff *et al.*, 1988, *Nature* 334, 585-591
- Heid et al., 1996, *Genome Research* 6: 986-994
- Holland et al., 1991, *PNAS* 88: 7276-7280
- Iwabuchi *et al.*, 1993, *Oncogene* 8, 1693-1696
- Jeffreys et al., 1985, *Nature* 316: 76-9
- 15 Johnson *et al.*, 1989, *Endoc. Rev.* 10, 317-331
- Kellogg et al., 1990, *Anal. Biochem.* 189:202-208
- Lam, 1997, *Anticancer Drug Res.* 12(3):145-67
- Lesage et al. (1996a), *Genomics* 34: 153-155
- Lesage et al. (1996b), *EMBO J.* 15: 1004-1011
- 20 Livak et al., 1995, *PCR Methods and Applications* 357-362
- Logan, Shenk, 1984, *Proc. Natl. Acad. Sci.* 81, 3655-3659
- Lowy *et al.*, 1980, *Cell* 22, 817-23
- Maddox *et al.*, 1983, *J. Exp. Med.* 158, 1211-1216
- McConnell *et al.*, 1992, *Science* 257, 1906-1912
- 25 Nicholls *et al.*, 1993, *J. Immunol. Meth.* 165, 81-91
- Piatak et al., 1993, *BioTechniques* 14:70-81
- Piatak et al., 1993, *Science* 259:1749-1754
- Porath *et al.*, 1992, *Prot. Exp. Purif.* 3, 263-281
- Roberge *et al.*, 1995, *Science* 269, 202-204
- 30 Schnitzler et al. 2003, *Receptor Channels* 9(1):79-85
- Scott & Smith 1990, *Science* 249:386-390
- Sjolander, Urbaniczky, 1991, *Anal. Chem.* 63, 2338-2345
- Szabo *et al.*, 1995, *Curr. Opin. Struct. Biol.* 5, 699-705
- Terstappen 1999: *Analyt. Biochem.* 272, 149-155
- 35 Thomas, 1980, *Proc. Nat. Acad. Sci.*, 77:5201-5205
- Uhlmann *et al.*, 1987, *Tetrahedron. Lett.* 215, 3539-3542

Weber et al., 1990, *Genomics* 7: 524-30

Wigler *et al.*, 1977, *Cell* 11, 223-32

Wigler *et al.*, 1980, *Proc. Natl. Acad. Sci.* 77, 3567-70

Willumsen et al. 2003, *Receptors and Ion Channels* 9: 3-12

Claims

1. A method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising the steps of
 - i) contacting a test compound with a KCNK1 polypeptide,
 - ii) detect binding of said test compound to said KCNK1 polypeptide.
2. A method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising the steps of
 - i) determining the functional activity of a KCNK1 polypeptide at a certain concentration of a test compound or in the absence of said test compound,
 - ii) determining the functional activity of said polypeptide at a different concentration of said test compound.
 - iii) comparing the functional activity of said polypeptide under conditions (i) and (ii) to the functional activity of the polypeptide in the presence of a known regulator.
3. The method according to claim 2 comprising
 - i) a cell expressing KCNK1 having functional activity
 - ii) contacting said cell with the test compounds
 - iii) measuring functional activity.
4. The method of claim 2 and 3 where the cell is expressing endogenous KCNK1.
5. The method of claim 2 and 3 where the cell is expressing recombinant KCNK1.
6. The method of claim 2 to 5 where the functional activity is the KCNK1 current.
7. The method of claim 2 to 5 where the functional activity is a change in membrane potential.
8. The method of claim 2 to 5 where the functional activity is a change in ion concentration.

9. The method of claim 2 to 5 where the functional activity is any other measurable effect exerted by KCNK1.
10. The method of any of claims 1 to 9, wherein the step of contacting is in or at the surface of a cell.
- 5 11. The method of any of claims 1 to 10, wherein the cell is in vitro.
12. The method of claim 1, wherein the step of contacting is in a cell-free system.
13. The method of any of claims 1 to 12, wherein the polypeptide is coupled to a detectable label.
14. The method of any of claims 1 to 13, wherein the compound is coupled to a detectable
10 label.
15. The method of any of claims 1 to 14, wherein the test compound displaces a ligand which is first bound to the polypeptide.
16. The method of claim 1, wherein the polypeptide is attached to a solid support.
17. The method of any of claims 1 to 9, wherein the compound is attached to a solid support.
- 15 18. A method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising the steps of
 - i) contacting a test compound with a KCNK1 polynucleotide,
 - 20 ii) detect binding of said test compound to said KCNK1 polynucleotide.
19. The method of claim 18 wherein the nucleic acid molecule is RNA.
20. The method of claim 18 wherein the contacting step is in or at the surface of a cell.
21. The method of claim 18 wherein the contacting step is in a cell-free system.
22. The method of claim 18 wherein polynucleotide is coupled to a detectable label.
- 25 23. The method of claim 18 wherein the test compound is coupled to a detectable label.

24. A method of diagnosing a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising the steps of
- 5 i) determining the amount of a KCNK1 polynucleotide in a sample taken from said mammal,
- ii) determining the amount of KCNK1 polynucleotide in healthy and/or diseased mammals.
25. A pharmaceutical composition for the treatment of a disease comprised in a group of
10 diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising a therapeutic agent which binds to a KCNK1 polypeptide.
26. A pharmaceutical composition for the treatment of a disease comprised in a group of
15 diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising a therapeutic agent which regulates the functional activity of a KCNK1 polypeptide.
27. A pharmaceutical composition for the treatment of a disease comprised in a group of
20 diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising a therapeutic agent which regulates the functional activity of a KCNK1 polypeptide, wherein said therapeutic agent is
- 25 i) a small molecule,
ii) an RNA molecule,
iii) an antisense oligonucleotide,
iv) a polypeptide,
v) an antibody, or
vi) a ribozyme.
- 30 28. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological

diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising a KCNK1 polynucleotide.

29. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising a KCNK1 polypeptide.
30. Use of regulators of a KCNK1 for the preparation of a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal.
31. Method for the preparation of a pharmaceutical composition useful for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal comprising the steps of
- i) identifying a regulator of KCNK1,
 - ii) determining whether said regulator ameliorates the symptoms of a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases in a mammal; and
 - iii) combining of said regulator with an acceptable pharmaceutical carrier.
32. Use of a regulator of KCNK1 for the regulation of KCNK1 functional activity in a mammal having a disease comprised in a group of diseases consisting of cardiovascular diseases, dermatological diseases, gastroenterological diseases, neurological diseases, hematological diseases, cancer, inflammation and urological diseases.

Fig. 1**SEQ ID NO: 1**

GGGCAGGAAGACGGCGCTGCCCGGAGGAGCGGGGCGGGCGGGCGCGCG
GGGGAGCGGGCGGCGGGCGGGAGCCAGGCCCGGGCGGGGGCGGGGGCG
GCGGGGCCAGAAGAGGCGGGCGGGCCGCGCTCCGGCCGGTCTGCGGCGT
TGGCCTTGGCTTTGGCTTTGGCGGCGGCGGTGGAGAAGATGCTGCAGT
CCCTGGCCGGCAGCTCGTGCGTGCGCCTGGTGGAGCGGCACCGCTCGG
CCTGGTGCTTCGGCTTCCTGGTGCTGGGCTACTTGCTCTACCTGGTCT
TCGGCGCAGTGGTCTTCTCCTCGGTGGAGCTGCCCTATGAGGACCTGC
TGCGCCAGGAGCTGCGCAAGCTGAAGCGACGCTTCTTGGAGGAGCACG
AGTGCCTGTCTGAGCAGCAGCTGGAGCAGTTCCTGGGCCGGGTGCTGG
AGGCCAGCAACTACGGCGTGTCGGTGCTCAGCAACGCCTCGGGCAACT
GGA ACTGGGACTTCACCTCCGCGCTCTTCTTCGCCAGCACCGTGCTCT
CCACCACAGGTTATGGCCACACCGTGCCCTTGTCAGATGGAGGTAAGG
CCTTCTGCATCATCTACTCCGTCATTGGCATTCCCTTCACCTCCTGT
TCCTGACGGCTGTGGTCCAGCGCATCACCGTGACGTCACCCGCAGGC
CGGTCCTCTACTTCCACATCCGCTGGGGCTTCTCCAAGCAGGTGGTGG
CCATCGTCCATGCCGTGCTCCTTGGGTTTGTCACTGTGTCCTGCTTCT
TCTTCATCCCGGCCGCTGTCTTCTCAGTCCTGGAGGATGACTGGA ACT
TCCTGGAATCCTTTTATTTTTTGTTTTATTTCCCTGAGCACCATTGGCC
TGGGGGATTATGTGCCTGGGGAAGGCTACAATCAAAAATTCAGAGAGC
TCTATAAGATTGGGATCACGTGTTACCTGCTACTTGGCCTTATTGCCA
TGTTGGTAGTTCTTGAAACCTTCTGTGAACTCCATGAGCTGAAAAAAT
TCAGAAAAATGTTCTATGTGAAGAAGGACAAGGACGAGGATCAGGTGC
ACATCATAGAGCATGACCAACTGTCCTTCTCCTCGATCACAGACCAGG
CAGCTGGCATGAAAGAGGACCAGAAGCAAAATGAGCCTTTTGTGGCCA
CCCAGTCATCTGCCTGCGTGGATGGCCCTGCAAACCATTGAGCGTAGG
ATTTGTTGCATTATGCTAGAGCACCAGGGTCAGGGTGCAAGGAAGAGG
CTTAAGTATGTTCA TTTTTATCAGAATGCAAAAGCGAAAATTATGTCA
CTTTAAGAAATAGCTACTGTTTGCAATGTCTTATTA AAAAACAACAAA
AAAAGACACATGGAACAAAGAAGCTGTGACCCCAGCAGGATGTCTAAT

Fig. 1 (continued)

ATGTGAGGAAATGAGATGTCCACCTAAAATTCATATGTGACAAAATTA
TCTCGACCTTACATAGGAGGAGAATACTTGAAGCAGTATGCTGCTGTG
GTTAGAAGCAGATTTTATACTTTTAACTGGAACTTTGGGGTTTGCAT
TTAGATCATTTAGCTGATGGCTAAATAGCAAATTTATATTTAGAAGC
AAAAAAAAAAGCATAGAGATGTGTTTTATAAATAGGTTTATGTGTAC
TGGTTTGCATGTACCCACCCAAAATGATTATTTTTGGAGAATCTAAGT
CAAAC TCACTATTTATAATGCATAGGTAACCATTA ACTATGTACATAT
AAAGTATAAATATGTTTATATTCTGTACATATGGTTTAGGTCACCAGA
TCCTAGTGTAGTTCTGAAACTAAGACTATAGATATTTTGTTCCTTTG
ATTTCTCTTTATACTAAAGAATCCAGAGTTGCTACAATAAAATAAGGG
GAATAATAAACTTGAGAGTGAATAACCAT

Fig. 2**SEQ ID NO: 2**

MLQSLAGSSCVRLVERHRSAWCFGFLVLGYLLYL VFGAVVFSSVELPY
EDLLRQELRKLKRRFLEEHECLSEQQLEQFLGRVLEASNYGVS VLSNA
SGNWNWDFTSALFFASTVLSTTGYGHTVPLSDGGKAF CIIYSVIGIPF
TLLFLTAVVQORITVHVTRRPVLYFHIRWGF SKQVVAIVHAVLLGFVTV
SCFFFIPAAVFSVLEDDWNFLESFYFCFISLSTIGLGDYVPGE GYNQK
FRELYKIGITCYLLLGLIAMLVLETFCELHELKKFRKMFYVKKDKDE
DQVHII EHDQLSFSSITDQAAGMKEDQKQNEPFVATQSSACVDGPANH

Fig. 3

SEQ ID NO: 3

5' - tgaagaaggacaaggacgagga - 3'

Fig. 4

SEQ ID NO: 4

5' - gcctggtctgtgatcgagga - 3'

Fig. 5

SEQ ID NO: 5

5' - caggtgcacatcatagagcatgaccaa - 3'

SEQUENCE LISTING

<110> Bayer HealthCare AG
 <120> Diagnostics and Therapeutics for Diseases Associated with Potassium Channel, Subfamily K, Member 1 (KCNK1)
 <130> BHC 03 01 019
 <160> 5
 <170> PatentIn version 3.1
 <210> 1
 <211> 1901
 <212> DNA
 <213> Homo sapiens
 <400> 1

```

gggcaggaag acggcgctgc ccggaggagc ggggcgggcg ggcgcgcggg ggagcgggcg      60
gcgggcggga gccagggccc ggcgggggcg ggggcgggcg ggccagaaga ggcgggcgggc      120
cgcgctccgg ccggtctgcg gcgttggcct tggctttggc tttggcgcg gcggtggaga      180
agatgctgca gtccctggcc ggcagctcgt gcgtgcgcct ggtggagcgg caccgctcgg      240
cctggtgctt cggcttcctg gtgctgggct acttgctcta cctggtcttc ggcgcagtg      300
tcttctcctc ggtggagctg ccctatgagg acctgctgcg ccaggagctg cgcaagctga      360
agcgacgctt cttggaggag cagcagtgcc tgtctgagca gcagctggag cagttcctgg      420
gccgggtgct ggaggccagc aactacggcg tgcgtgtgct cagcaacgcc tcggggcaact      480
ggaaactggga cttcacctcc gcgctcttct tcgccagcac cgtgctctcc accacagggt      540
atggccacac cgtgcccttg tcagatggag gtaaggcctt ctgcatcatc tactccgtca      600
ttggcattcc cttcacctcc ctgttcctga cggctgtggc ccagcgcatc accgtgcacg      660
tcacccgcag gccggtcctc tacttccaca tccgctgggg cttctccaag cagggtggtg      720
ccatcgtcca tgccgtgctc cttgggtttg tcaactgtgc ctgcttcttc ttcacccgg      780
ccgctgtctt ctcagtcctg gaggatgact ggaacttcct ggaatccttt tatttttgtt      840
ttatttccct gagcaccatt ggcttggggg attatgtgcc tggggaaggc tacaatcaaa      900
aattcagaga gctctataag attgggatca cgtgttacct gctacttggc cttattgcca      960
tgttggtagt tctggaaacc ttctgtgaac tccatgagct gaaaaaattc agaaaaatgt      1020
tctatgtgaa gaaggacaag gacgaggatc aggtgcacat catagagcat gaccaactgt      1080
ccttctcctc gatcacagac caggcagctg gcatgaaaga ggaccagaag caaaatgagc      1140
cttttgtggc caccagtc tctgcctgcg tggatggccc tgcaaaacct tgagcgtagg      1200
atttggtgca ttatgctaga gcaccagggt cagggtgcaa ggaagaggct taagtatgtt      1260
cattttttatc agaatgcaaa agcgaataat atgtcacttt aagaaatagc tactgtttgc      1320
aatgtcttat taaaaaaca caaaaaaga cacatggaac aaagaagctg tgaccccagc      1380
aggatgtcta atatgtgagg aaatgagatg tccacctaaa attcatatgt gacaaaatta      1440
tctcgacctt acataggagg agaatacttg aagcagtatg ctgctgtggt tagaagcaga      1500
ttttatactt ttaactggaa actttggggg ttgcatttag atcatttagc tgatggctaa      1560
atagcaaaat ttatathtag aagcaaaaaa aaaaagcata gagatgtggt ttataaatag      1620
gtttatgtgt actggtttgc atgtaccac ccaaaatgat tatttttga gaatctaagt      1680
caaactcact atttataatg cataggtaac cattaactat gtacatataa agtataaata      1740
tgtttatatt ctgtacatat ggttttaggtc accagatcct agttagttc tgaaactaag      1800
actatagata tttgtttct tttgatttct ctttatacta aagaatccag agttgctaca      1860
ataaaataag gggaataata aacttgagag tgaataacca t      1901

```

<210> 2
 <211> 336
 <212> PRT
 <213> Homo sapiens
 <400> 2

```

Met Leu Gln Ser Leu Ala Gly Ser Ser Cys Val Arg Leu Val Glu Arg
1          5          10          15
His Arg Ser Ala Trp Cys Phe Gly Phe Leu Val Leu Gly Tyr Leu Leu
20          25          30
Tyr Leu Val Phe Gly Ala Val Val Phe Ser Ser Val Glu Leu Pro Tyr
35          40          45
Glu Asp Leu Leu Arg Gln Glu Leu Arg Lys Leu Lys Arg Arg Phe Leu
50          55          60
Glu Glu His Glu Cys Leu Ser Glu Gln Gln Leu Glu Gln Phe Leu Gly
65          70          75          80
Arg Val Leu Glu Ala Ser Asn Tyr Gly Val Ser Val Leu Ser Asn Ala
85          90          95
Ser Gly Asn Trp Asn Trp Asp Phe Thr Ser Ala Leu Phe Phe Ala Ser
100         105         110
Thr Val Leu Ser Thr Thr Gly Tyr Gly His Thr Val Pro Leu Ser Asp
115         120         125
Gly Gly Lys Ala Phe Cys Ile Ile Tyr Ser Val Ile Gly Ile Pro Phe
130         135         140
Thr Leu Leu Phe Leu Thr Ala Val Val Gln Arg Ile Thr Val His Val
145         150         155         160

```

- 2 -

Thr Arg Arg Pro Val Leu Tyr Phe His Ile Arg Trp Gly Phe Ser Lys
 165 170 175
 Gln Val Val Ala Ile Val His Ala Val Leu Leu Gly Phe Val Thr Val
 180 185 190
 Ser Cys Phe Phe Phe Ile Pro Ala Ala Val Phe Ser Val Leu Glu Asp
 195 200 205
 Asp Trp Asn Phe Leu Glu Ser Phe Tyr Phe Cys Phe Ile Ser Leu Ser
 210 215 220
 Thr Ile Gly Leu Gly Asp Tyr Val Pro Gly Glu Gly Tyr Asn Gln Lys
 225 230 235 240
 Phe Arg Glu Leu Tyr Lys Ile Gly Ile Thr Cys Tyr Leu Leu Leu Gly
 245 250 255
 Leu Ile Ala Met Leu Val Val Leu Glu Thr Phe Cys Glu Leu His Glu
 260 265 270
 Leu Lys Lys Phe Arg Lys Met Phe Tyr Val Lys Lys Asp Lys Asp Glu
 275 280 285
 Asp Gln Val His Ile Ile Glu His Asp Gln Leu Ser Phe Ser Ser Ile
 290 295 300
 Thr Asp Gln Ala Ala Gly Met Lys Glu Asp Gln Lys Gln Asn Glu Pro
 305 310 315 320
 Phe Val Ala Thr Gln Ser Ser Ala Cys Val Asp Gly Pro Ala Asn His
 325 330 335

<210> 3
 <211> 22
 <212> DNA
 <213> artificial sequence
 <220>
 <223> forward primer

<400> 3
 tgaagaagga caaggacgag ga

22

<210> 4
 <211> 20
 <212> DNA
 <213> artificial sequence
 <220>
 <223> reverse primer
 <400> 4
 gcctggtctg tgatcgagga

20

<210> 5
 <211> 27
 <212> DNA
 <213> artificial sequence
 <220>
 <223> probe
 <400> 5
 caggtgcaca tcatagagca tgaccaa

27